


512.94

M55s

1906

cop. 2





LIBRARY
OF THE
UNIVERSITY
OF ILLINOIS

PRESENTED BY
PROF. E. G. LEWIS
FROM THE LIBRARY OF
M. W. JACOBS, JR.

1950

512.2 512.94
M55s4

cop. 2

MATHEMATICS LIBRARY

Return this book on or before the
Latest Date stamped below.

University of Illinois Library

MAR 19 1964

APR 13 1964

MAY 25 1964

AUG 8 1964

SEP 10 1964

SEP 1 1967
AUG 7 REC'D

MAY 04 1991

APR 22 REC'D

JUL 27 REC'D

OCT 20 1998

JUL 09 REC'D

MATHEMATICAL MONOGRAPHS.

EDITED BY

Mansfield Merriman and Robert S. Woodward.

Octavo, Cloth, \$1.00 each.

No. 1. HISTORY OF MODERN MATHEMATICS.

By DAVID EUGENE SMITH.

No. 2. SYNTHETIC PROJECTIVE GEOMETRY.

By GEORGE BRUCE HALSTED.

No. 3. DETERMINANTS.

By LAENAS GIFFORD WELD.

No. 4. HYPERBOLIC FUNCTIONS.

By JAMES McMAHON.

No. 5. HARMONIC FUNCTIONS.

By WILLIAM E. BYERLY.

No. 6. GRASSMANN'S SPACE ANALYSIS.

By EDWARD W. HYDE.

No. 7. PROBABILITY AND THEORY OF ERRORS.

By ROBERT S. WOODWARD.

No. 8. VECTOR ANALYSIS AND QUATERNIONS.

By ALEXANDER MACFARLANE.

No. 9. DIFFERENTIAL EQUATIONS.

By WILLIAM WOOLSEY JOHNSON.

No. 10. THE SOLUTION OF EQUATIONS.

By MANSFIELD MERRIMAN.

No. 11. FUNCTIONS OF A COMPLEX VARIABLE.

By THOMAS S. FISKE.

PUBLISHED BY

JOHN WILEY & SONS, NEW YORK.

CHAPMAN & HALL, Limited, LONDON.

MATHEMATICAL MONOGRAPHS.

EDITED BY

MANSFIELD MERRIMAN AND ROBERT S. WOODWARD.

No. 10.

THE SOLUTION OF
EQUATIONS.

BY

MANSFIELD MERRIMAN,

PROFESSOR OF CIVIL ENGINEERING IN LEHIGH UNIVERSITY.

FOURTH EDITION, ENLARGED.

FIRST THOUSAND.

NEW YORK:

JOHN WILEY & SONS.

LONDON: CHAPMAN & HALL, LIMITED.

1906.

COPYRIGHT, 1896,
BY
MANSFIELD MERRIMAN AND ROBERT S. WOODWARD
UNDER THE TITLE
HIGHER MATHEMATICS.

First Edition, September, 1896.

Second Edition, January, 1898.

Third Edition, August, 1900.

Fourth Edition, January, 1906.

ROBERT DRUMMOND, PRINTER, NEW YORK.

EDITORS' PREFACE.

THE volume called Higher Mathematics, the first edition of which was published in 1896, contained eleven chapters by eleven authors, each chapter being independent of the others, but all supposing the reader to have at least a mathematical training equivalent to that given in classical and engineering colleges. The publication of that volume is now discontinued and the chapters are issued in separate form. In these reissues it will generally be found that the monographs are enlarged by additional articles or appendices which either amplify the former presentation or record recent advances. This plan of publication has been arranged in order to meet the demand of teachers and the convenience of classes, but it is also thought that it may prove advantageous to readers in special lines of mathematical literature.

It is the intention of the publishers and editors to add other monographs to the series from time to time, if the call for the same seems to warrant it. Among the topics which are under consideration are those of elliptic functions, the theory of numbers, the group theory, the calculus of variations, and non-Euclidean geometry; possibly also monographs on branches of astronomy, mechanics, and mathematical physics may be included. It is the hope of the editors that this form of publication may tend to promote mathematical study and research over a wider field than that which the former volume has occupied.

December, 1905.

AUTHOR'S PREFACE.

THE following pages are designed as supplementary to the discussions of equations in college text-books, and several methods of solution not commonly given in such works are presented and exemplified. The aim kept in view has been that of the determination of the numerical values of the roots of numerical equations, and algebraic analysis has been used only to further this end. Historical references are given, problems stated as exercises for the student, and the attempt has everywhere been made to present the subject clearly and concisely. The volume has not been written for those thoroughly conversant with the theory of equations, but rather for students of mathematics, computers, and engineers.

This edition has been enlarged by the addition of five articles which render the former treatment more complete and also give recent investigations regarding the expression of roots in series. While not designed for college classes, it is hoped that the book may prove useful to postgraduate students in mathematics, physics and engineering, and also tend to promote general interest in mathematical science.

SOUTH BETHLEHEM, PA.,
December, 1905.

CONTENTS.

ART. 1. INTRODUCTION	Page	1
2. GRAPHIC SOLUTIONS	3	
3. THE REGULA FALSI	5	
4. NEWTON'S APPROXIMATION RULE	6	
5. SEPARATION OF THE ROOTS	8	
6. NUMERICAL ALGEBRAIC EQUATIONS	10	
7. TRANSCENDENTAL EQUATIONS	13	
8. ALGEBRAIC SOLUTIONS	15	
9. THE CUBIC EQUATION	17	
10. THE QUARTIC EQUATION	19	
11. QUINTIC EQUATIONS	21	
12. TRIGONOMETRIC SOLUTIONS	24	
13. REAL ROOTS BY SERIES	27	
14. COMPUTATION OF ALL ROOTS	28	
15. ROOTS OF UNITY	31	
16. SOLUTIONS BY MACLAURIN'S SERIES	33	
17. SYMMETRIC FUNCTIONS OF ROOTS	37	
18. LOGARITHMIC SOLUTIONS	39	
19. INFINITE EQUATIONS	43	
20. NOTES AND PROBLEMS	45	
INDEX	47	

THE SOLUTION OF EQUATIONS.

ART. 1. INTRODUCTION.

THE science of algebra arose in the efforts to solve equations. Indeed algebra may be called the science of the equation, since the discussion of equalities and the transformation of forms into simpler equivalent ones have been its main objects. The solution of an equation containing one unknown quantity consists in the determination of its value or values, these being called roots. An algebraic equation of degree n has n roots, while transcendental equations often have an infinite number of roots. The object of the following pages is to present and exemplify convenient methods for the determination of the numerical values of the roots of both kinds of equations, the real roots receiving special attention because these are mainly required in the solution of problems in physical science.

An algebraic equation is one that involves only the operations of arithmetic. It is to be first freed from radicals so as to make the exponents of the unknown quantity all integers; the degree of the equation is then indicated by the highest exponent of the unknown quantity. The algebraic solution of an algebraic equation is the expression of its roots in terms of the literal coefficients; this is possible, in general, only for linear, quadratic, cubic, and quartic equations, that is, for equations of the first, second, third, and fourth degrees. A numerical equation is an algebraic equation having all its coefficients real numbers, either positive or negative. For the four degrees

above mentioned the roots of numerical equations may be computed from the formulas for the algebraic solutions, unless they fall under the so-called irreducible case wherein real quantities are expressed in imaginary forms.

An algebraic equation of the n^{th} degree may be written with all its terms transposed to the first member, thus:

$$x^n + a_1x^{n-1} + a_2x^{n-2} + \dots + a_{n-1}x + a_n = 0,$$

and, for brevity, the first member will be called $f(x)$ and the equation be referred to as $f(x) = 0$. The roots of this equation are the values of x which satisfy it, that is, those values of x that reduce $f(x)$ to 0. When all the coefficients a_1, a_2, \dots, a_n are real, as will always be supposed to be the case, Sturm's theorem gives the number of real roots, provided they are unequal, as also the number of real roots lying between two assumed values of x , while Horner's method furnishes a convenient process for obtaining the values of the roots to any required degree of precision.

A transcendental equation is one involving the operations of trigonometry or of logarithms, as, for example, $x + \cos x = 0$, or $a^{2x} + xb^x = 0$. No general method for the literal solution of these equations exists; but when all known quantities are expressed as real numbers, the real roots may be located and computed by tentative methods. Here also the equation may be designated as $f(x) = 0$, and the discussions in Arts. 2-5 will apply equally well to both algebraic and transcendental forms. The methods to be given are thus, in a sense, more valuable than Sturm's theorem and Horner's process, although for algebraic equations they may be somewhat longer. It should be remembered, however, that algebraic equations higher than the fourth degree do not often occur in physical problems, and that the value of a method of solution is to be measured not merely by the rapidity of computation, but also by the ease with which it can be kept in mind and applied.

Prob. 1. Reduce the equation $(a + x)^{\frac{1}{2}} + (a - x)^{\frac{1}{2}} = 2b$ to an equation having the exponents of the unknown quantity all integers.


ART. 2. GRAPHIC SOLUTIONS.

Approximate values of the real roots of two simultaneous algebraic equations may be found by the methods of plane analytic geometry when the coefficients are numerically expressed. For example, let the given equations be

$$x^2 + y^2 = a^2, \quad x^2 - bx = y^2 - cy,$$

the first representing a circle and the second a hyperbola. Drawing two rectangular axes OX and OY , the circle is described from O with the radius a . The coordinates of the center of the hyperbola are found to be $OA = \frac{1}{2}b$ and $AC = \frac{1}{2}c$, while its diameter $BD = \sqrt{b^2 - c^2}$, from which the two branches may be described.

The intersections of the circle with the hyperbola give the real values of x and y . If $a = 1$, $b = 4$, and $c = 3$, there are but two real values for x and two real values for y , since the circle intersects but one branch of the hyperbola; here Om is the positive and




Op the negative value of x , while mn is the positive and pq the negative value of y . When the radius a is so large that the circle intersects both branches of the hyperbola there are four real values of both x and y .

By a similar method approximate values of the real roots of an algebraic equation containing but one unknown quantity may be graphically found. For instance, let the cubic equation $x^3 + ax - b = 0$ be required to be solved.* This may be written as the two simultaneous equations

$$y = x^3, \quad y = -ax + b,$$

* See Proceedings of the Engineers' Club of Philadelphia, 1884, Vol. IV, pp. 47-49

and the graph of each being plotted, the abscissas of their points of intersection give the real roots of the cubic. The



curve $y = x^3$ should be plotted upon cross-section paper by the help of a table of cubes; then OB is laid off equal to b , and OC equal to a/b , taking care to observe the signs of a and b . The line joining B and C cuts the curve at p , and hence qp is the real root of $x^3 + ax - b = 0$. If the

cubic equation have three real roots the straight line BC will intersect the curve in three points.

Some algebraic equations of higher degrees may be graphically solved in a similar manner. For the quartic equation $x^4 + Ax^2 + Bx - C = 0$, it is best to put $z = A^{\frac{1}{2}}x$, and thus reduce it to the form $x^4 + x^2 + bx - c = 0$; then the two equations to be plotted are

$$y = x^4 + x^2, \quad y = -bx + c,$$

the first of which may be drawn once for all upon cross-section paper, while the straight line represented by the second may be drawn for each particular case, as described above.*

This method is also applicable to many transcendental equations; thus for the equation $Ax - B\sin x = 0$ it is best to write $ax - \sin x = 0$; then $y = \sin x$ is readily plotted by help of a table of sines, while $y = ax$ is a straight line passing through the origin. In the same way $a^x - x^2 = 0$ gives the curve represented by $y = a^x$ and the parabola represented by $y = x^2$, the intersections of which determine the real roots of the given equation.

Prob. 2. Devise a graphic solution for finding approximate values of the real roots of the equation $x^5 + ax^3 + bx^2 + cx + d = 0$.

Prob. 3. Determine graphically the number and the approximate values of the real roots of the equation $\arcsin x - 8 \sin x = 0$. (Ans.—Six real roots, $x = \pm 159^\circ$, $\pm 430^\circ$, and $\pm 456^\circ$.)


* For an extension of this method to the determination of imaginary roots, see Phillips and Beebe's *Graphic Algebra*, New York, 1882.

ART. 3. THE REGULA FALSI.

One of the oldest methods for computing the real root of an equation is the rule known as "regula falsi," often called the method of double position.* It depends upon the principle that if two numbers x_1 and x_2 be substituted in the expression $f(x)$, and if one of these renders $f(x)$ positive and the other renders it negative, then at least one real root of the equation $f(x) = 0$ lies between x_1 and x_2 . Let the figure represent a part of the real graph of the equation $y = f(x)$. The point X , where the curve crosses the axis of abscissas, gives a real root OX of the equation $f(x) = 0$. Let OA and OB be inferior and superior limits of the root OX which are determined either by trial or by the method of Art. 5.

Let Aa and Bb be the values of $f(x)$ corresponding to these limits.

Join ab , then the intersection C of the straight line ab with the axis OB gives an approximate value OC for the root. Now compute



Cc and join ac , then the intersection D gives a value OD which is closer still to the root OX .

Let x_1 and x_2 be the assumed values OA and OB , and let $f(x_1)$ and $f(x_2)$ be the corresponding values of $f(x)$ represented by Aa and Bb , these values being with contrary signs. Then from the similar triangle AaC and BbC the abscissa OC is

$$x_3 = \frac{x_2 f(x_1) - x_1 f(x_2)}{f(x_1) - f(x_2)} = x_1 + \frac{(x_2 - x_1) f(x_1)}{f(x_1) - f(x_2)} = x_2 + \frac{(x_2 - x_1) f(x_2)}{f(x_1) - f(x_2)}.$$

By a second application of the rule to x_1 and x_3 , another value x_4 is computed, and by continuing the process the value of x can be obtained to any required degree of precision.

As an example let $f(x) = x^5 + 5x^2 + 7 = 0$. Here it may be found by trial that a real root lies between -2 and -1.8 .

* This originated in India, and its first publication in Europe was by Abraham ben Esra, in 1130. See Matthiesen, Grundzüge der antiken und modernen Algebra der literalen Gleichungen, Leipzig, 1878.

For $x_1 = -2$, $f(x_1) = -5$, and for $x_2 = -1.8$, $f(x_2) = +4.304$; then by the regula falsi there is found $x_3 = -1.90$ nearly. Again, for $x_3 = -1.90$, $f(x_3) = +0.290$, and these combined with x_1 and $f(x_1)$ give $x_4 = -1.906$, which is correct to the third decimal.


As a second example let $f(x) = \arcsin x - \sin x - 0.5 = 0$. Here a graphic solution shows that there is but one real root, and that the value of it lies between 85° and 86° . For $x_1 = 85^\circ$, $f(x_1) = -0.01266$, and for $x_2 = 86^\circ$, $f(x_2) = +0.00342$; then by the rule $x_3 = 85^\circ 44'$, which gives $f(x_3) = -0.00090$. Again, combining the values for x_2 and x_3 there is found $x_4 = 85^\circ 47'$, which gives $f(x_4) = -0.00009$. Lastly, combining the values for x_2 and x_4 there is found $x_5 = 85^\circ 47'.4$, which is as close an approximation as can be made with five-place tables.

In the application of this method it is to be observed that the signs of the values of x and $f(x)$ are to be carefully regarded, and also that the values of $f(x)$ to be combined in one operation should have opposite signs. For the quickest approximation the values of $f(x)$ to be selected should be those having the smallest numerical values.

Prob. 4. Compute by the regula falsi the real roots of $x^5 - 0.25 = 0$. Also those of $x^2 + \sin 2x = 0$.

ART. 4. NEWTON'S APPROXIMATION RULE.

Another useful method for approximating to the value of the real root of an equation is that devised by Newton in 1666.*



If $y = f(x)$ be the equation of a curve, OX in the figure represents a real root of the equation $f(x) = 0$. Let OA be an approximate value of OX , and Aa the corresponding value of $f(x)$. At a let aB be drawn tangent to the curve; then OB is another approximate value of OX .

* See Analysis per equationes numero terminorum infinitas, p. 269, Vol. I of Horsely's edition of Newton's works (London, 1779), where the method is given in a somewhat different form.

Let Bb be the value of $f(x)$ corresponding to OB , and at b let the tangent bC be drawn; then OC is a closer approximation to OX , and thus the process may be continued.

Let $f'(x)$ be the first derivative of $f(x)$; or, $f'(x) = df(x)/dx$. For $x = x_1 = OA$ in the figure, the value of $f(x_1)$ is the ordinate Aa , and the value of $f'(x_1)$ is the tangent of the angle aBA ; this tangent is also Aa/AB . Hence $AB = f(x_1)/f'(x_1)$, and accordingly OB and OC are found by

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}, \quad x_3 = x_2 - \frac{f(x_2)}{f'(x_2)},$$

which is Newton's approximation rule. By a third application to x_3 the closer value x_4 is found, and the process may be continued to any degree of precision required.

For example, let $f(x) = x^5 + 5x^2 + 7 = 0$. The first derivative is $f'(x) = 5x^4 + 10x$. Here it may be found by trial that -2 is an approximate value of the real root. For $x_1 = -2$ $f(x_1) = -5$, and $f'(x_1) = 60$, whence by the rule $x_2 = -1.92$. Now for $x_2 = -1.92$ are found $f(x_2) = -0.6599$ and $f'(x_2) = 29.052$, whence by the rule $x_3 = -1.906$, which is correct to the third decimal.

As a second example let $f(x) = x^2 + 4 \sin x = 0$. Here the first derivative is $f'(x) = 2x + 4 \cos x$. An approximate value of x found either by trial or by a graphic solution is $x = -1.94$, corresponding to about $-111^\circ 09'$. For $x_1 = -1.94$, $f(x_1) = 0.03304$ and $f'(x_1) = -5.323$, whence by the rule $x_2 = -1.934$. By a second application $x_3 = -1.9328$, which corresponds to an angle of $-110^\circ 54\frac{1}{2}'$.

In the application of Newton's rule it is best that the assumed value of x_1 should be such as to render $f(x_1)$ as small as possible, and also $f'(x_1)$ as large as possible. The method will fail if the curve has a maximum or minimum between a and b . It is seen that Newton's rule, like the regula falsi, applies equally well to both transcendental and algebraic equations, and moreover that the rule itself is readily kept in mind by help of the diagram.

Prob. 5. Compute by Newton's rule the real roots of the algebraic equation $x^4 - 7x + 6 = 0$. Also the real roots of the transcendental equation $\sin x + \arcsin x - 2 = 0$.

ART. 5. SEPARATION OF THE ROOTS.


The roots of an equation are of two kinds, real roots and imaginary roots. Equal real roots may be regarded as a special class, which lie at the limit between the real and the imaginary. If an equation has p equal roots of one value and q equal roots of another value, then its first derivative equation has $p - 1$ roots of the first value and $q - 1$ roots of the second value, and thus all the equal roots are contained in a factor common to both primitive and derivative. Equal roots may hence always be readily detected and removed from the given equation. For instance, let $x^4 - 9x^2 + 4x + 12 = 0$, of which the derivative equation is $4x^3 - 18x + 4 = 0$; as $x - 2$ is a factor of these two equations, two of the roots of the primitive equation are ± 2 .

The problem of determining the number of the real and imaginary roots of an algebraic equation is completely solved by Sturm's theorem. If, then, two values be assigned to x the number of real roots between those limits is found by the same theorem, and thus by a sufficient number of assumptions limits may be found for each real root. As Sturm's theorem is known to all who read these pages, no applications of it will be here given, but instead an older method due to Hudde will be presented which has the merit of giving a comprehensive view of the subject, and which moreover applies to transcendental as well as to algebraic equations.*

If any equation $y = f(x)$ be plotted with values of x as abscissas and values of y as ordinates, a real graph is obtained whose intersections with the axis OX give the real roots of the

* Devised by Hudde in 1659 and published by Rolle in 1690. See *Œuvres de Lagrange*, Vol. VIII, p. 190.

equation $f(x) = 0$. Thus in the figure the three points marked X give three values OX for three real roots. The curve which represents $y = f(x)$ has points of maxima and minima marked A , and inflection points marked B . Now let the first deriva-



tive equation $dy/dx = f'(x)$ be formed and be plotted in the same manner on the axis $O'X'$. The condition $f'(x) = 0$ gives the abscissas of the points A , and thus the real roots $O'X'$ give limits separating the real roots of $f(x) = 0$. To ascertain if a real root OX lies between two values of $O'X'$ these two values are to be substituted in $f(x)$: if the signs of $f(x)$ are unlike in the two cases, a real root of $f(x) = 0$ lies between the two limits; if the signs are the same, a real root does not lie between those limits.

In like manner if the second derivative equation, that is, $d^2y/dx^2 = f''(x)$, be plotted on $O''X''$, the intersections give limits which separate the real roots of $f'(x) = 0$. It is also seen that the roots of the second derivative equation are the abscissas of the points of inflection of the curve $y = f(x)$.

To illustrate this method let the given equation be the quintic $f(x) = x^5 - 5x^3 + 6x + 2 = 0$. The first derivative equation is $f'(x) = 5x^4 - 15x^2 + 6 = 0$, the roots of which are approximately $-1.59, -0.69, +0.69, +1.59$. Now let each of these values be substituted for x in the given quintic, as also the values $-\infty, 0$, and $+\infty$, and let the corresponding values of $f(x)$ be determined as follows:

$$\begin{array}{l} x = -\infty, \quad -1.59, \quad -0.69, \quad 0, \quad +0.69, \quad +1.59, \quad +\infty; \\ f(x) = -\infty, \quad +2.4, \quad -0.6, \quad +2, \quad +4.7, \quad +1.6, \quad +\infty. \end{array}$$

Since $f(x)$ changes sign between $x_0 = -\infty$ and $x_1 = -1.59$, one real root lies between these limits; since $f(x)$ changes sign between $x_1 = -1.59$ and $x_2 = -0.69$, one real root lies between these limits; since $f(x)$ changes sign between $x_2 = -0.69$ and $x_3 = 0$, one real root lies between these limits; since $f(x)$ does not change sign between $x_3 = 0$ and $x_4 = \infty$, a pair of imaginary roots is indicated, the sum of which lies between $+0.69$ and ∞ .

As a second example let $f(x) = e^x - e^{2x} - 4 = 0$. The first derivative equation is $f'(x) = e^x - 2e^{2x} = 0$, which has two roots $e^x = \frac{1}{2}$ and $e^x = 0$, the latter corresponding to $x = -\infty$. For $x = -\infty$, $f(x)$ is negative; for $e^x = \frac{1}{2}$, $f(x)$ is negative; for $x = +\infty$, $f(x)$ is negative. The equation $e^x - e^{2x} - 4 = 0$ has, therefore, no real roots.

When the first derivative equation is not easily solved, the second, third, and following derivatives may be taken until an equation is found whose roots may be obtained. Then, by working backward, limits may be found in succession for the roots of the derivative equations until finally those of the primitive are ascertained. In many cases, it is true, this process may prove lengthy and difficult, and in some it may fail entirely; nevertheless the method is one of great theoretical and practical value.

Prob. 6. Show that $e^x + e^{-3x} - 4 = 0$ has two real roots, one positive and one negative.

Prob. 7. Show that $x^6 + x + 1 = 0$ has no real roots; also that $x^6 - x - 1 = 0$ has two real roots, one positive and one negative.

ART. 6. NUMERICAL ALGEBRAIC EQUATIONS.

An algebraic equation of the n^{th} degree may be written with all its terms transposed to the first member, thus:

$$x^n + a_1 x^{n-1} + a_2 x^{n-2} + \dots + a_{n-1} x + a_n = 0;$$

and if all the coefficients and the absolute term are real numbers, this is commonly called a numerical equation. The first member may for brevity be denoted by $f(x)$ and the equation itself by $f(x) = 0$.

The following principles of the theory of algebraic equations with real coefficients, deduced in text-books on algebra, are here recapitulated for convenience of reference :

(1) If x_1 is a root of the equation, $f(x)$ is divisible by $x - x_1$; and conversely, if $f(x)$ is divisible by $x - x_1$, then x_1 is a root of the equation.

(2) An equation of the n^{th} degree has n roots and no more.

(3) If x_1, x_2, \dots, x_n are the roots of the equation, then the product $(x - x_1)(x - x_2) \dots (x - x_n)$ is equal to $f(x)$.

(4) The sum of the roots is equal to $-a_1$; the sum of the products of the roots, taken two in a set, is equal to $+a_2$; the sum of the products of the roots, taken three in a set, is equal to $-a_3$; and so on. The product of all the roots is equal to $-a_n$ when n is odd, and to $+a_n$ when n is even.

(5) The equation $f(x) = 0$ may be reduced to an equation lacking its second term by substituting $y - a_1/n$ for x .*

(6) If an equation has imaginary roots, they occur in pairs of the form $p \pm qi$ where i represents $\sqrt{-1}$.

(7) An equation of odd degree has at least one real root whose sign is opposite to that of a_n .

(8) An equation of even degree, having a_n negative, has at least two real roots, one being positive and the other negative.

(9) A complete equation cannot have more positive roots than variations in the signs of its terms, nor more negative roots than permanences in signs. If all roots be real, there are as many positive roots as variations, and as many negative roots as permanences.†

(10) In an incomplete equation, if an even number of terms, say $2m$, are lacking between two other terms, then it has at least $2m$

* By substituting $y^2 + py + q$ for x , the quantities p and q may be determined so as to remove the second and third terms by means of a quadratic equation, the second and fourth terms by means of a cubic equation, or the second and fifth terms by means of a quartic equation.

† The law deduced by Harriot in 1631 and by Descartes in 1639.

imaginary roots; if an odd number of terms, say $2m + 1$, are lacking between two other terms, then it has at least either $2m + 2$ or $2m$ imaginary roots, according as the two terms have like or unlike signs.*

(11) Sturm's theorem gives the number of real roots, provided that they are unequal, as also the number of real roots lying between two assumed values of x .

(12) If a_r is the greatest negative coefficient, and if a_s is the greatest negative coefficient after x is changed into $-x$, then all real roots lie between the limits $a_r + 1$ and $-(a_s + 1)$.

(13) If a_h is the first negative and a_r the greatest negative coefficient, then $a_r^{\frac{1}{h}} + 1$ is a superior limit of the positive roots. If a_k be the first negative and a_s the greatest negative coefficient after x is changed into $-x$, then $a_s^{\frac{1}{k}} + 1$ is a numerically superior limit of the negative roots.

(14) Inferior limits of the positive and negative roots may be found by placing $x = z^{-1}$ and thus obtaining an equation $f(z) = 0$ whose roots are the reciprocals of $f(x) = 0$.

(15) Horner's method, using the substitution $x = z - r$ where r is an approximate value of x_1 , enables the real root x_1 to be computed to any required degree of precision.

The application of these principles and methods will be familiar to all who read these pages. Horner's method may be also modified so as to apply to the computation of imaginary roots after their approximate values have been found.† The older method of Hudde and Rolle, set forth in Art. 5, is however one of frequent convenient application, for such algebraic equations as actually arise in practice. By its use, together with principles (13) and (14) above, and the regula falsi of Art. 3, the real roots may be computed without any assumptions whatever regarding their values.

For example, let a sphere of diameter D and specific gravity

* Established by Du Gua; see Memoirs Paris Academy, 1741, pp. 435-494.

† Sheffler, Die Auflösung der algebraischen und transzendenten Gleichungen, Braunschweig, 1859; and Jelink, Die Auflösung der höheren numerischen Gleichungen, Leipzig, 1865.

g float in water, and let it be required to find the depth of immersion. The solution of the problem gives for the depth x the cubic equation

$$x^3 - \frac{3}{2}Dx^2 + \frac{1}{2}D^3g = 0.$$

As a particular case let $D = 2$ feet and $g = 0.65$; then the equation

$$x^3 - 3x^2 + 2.6 = 0$$

is to be solved. The first derivative equation is $3x^2 - 6x = 0$ whose roots are 0 and 2. Substituting these, there is found one negative root, one positive root less than 2, and one positive root greater than 2. The physical aspect of the question excludes the first and last root, and the second is to be computed. By (13) and (14) an inferior limit of this root is about 0.5, so that it lies between 0.5 and 2. For $x_1 = 0.5$, $f(x_1) = +1.975$, and for $x_2 = 2$, $f(x_2) = -1.4$; then by the regula falsi $x_3 = 1.35$. For $x_3 = 1.35$, $f(x_3) = -0.408$, and combining this with x , the regula falsi gives $x_4 = 1.204$ feet, which, except in the last decimal, is the correct depth of immersion of the sphere.

Prob. 8. The diameter of a water-pipe whose length is 200 feet and which is to discharge 100 cubic feet per second under a head of 10 feet is given by the real root of the quintic equation $x^5 - 38x - 101 = 0$. Find the value of x .

ART. 7. TRANSCENDENTAL EQUATIONS.

Rules (1) to (15) of the last article have no application to trigonometrical or exponential equations, but the general principles and methods of Arts. 2-5 may be always used in attempting their solution. Transcendental equations may have one, many, or no real roots, but those arising from problems in physical science must have at least one real root. Two examples of such equations will be presented.

A cylinder of specific gravity g floats in water, and it is required to find the immersed arc of the circumference. If this be expressed in circular measure it is given by the transcendental equation

$$f(x) = x - \sin x - 2\pi g = 0.$$

The first derivative equation is $1 - \cos x = 0$, whose root is any even multiple of 2π . Substituting such multiples in $f(x)$ it is found that the equation has but one real root, and that this lies between 0 and 2π ; substituting $\frac{1}{2}\pi$, $\frac{3}{4}\pi$, and π for x , it is further found that this root lies between $\frac{3}{4}\pi$ and π .

As a particular case let $g = 0.424$, and for convenience in using the tables let x be expressed in degrees; then

$$f(x) = x - 57^\circ.2958 \sin x - 152^\circ.64.$$

Now proceeding by the regula falsi (Art. 3) let $x_1 = 180^\circ$ and $x_2 = 135^\circ$, giving $f(x_1) = +27^\circ.36$ and $f(x_2) = -58^\circ.16$, whence $x_3 = 166^\circ$. For $x_3 = 166^\circ$, $f(x_3) = -0^\circ.469$, and hence 166° is an approximate value of the root. Continuing the process, x is found to be $166^\circ.237$, or in circular measure $x = 2.9014$ radians.

As a second example let it be required to find the horizontal tension of a catenary cable whose length is 22 feet, span 20 feet, and weight 10 pounds per linear foot, the ends being suspended from two points on the same level. If l be the span, s the length of the cable, and z a length of the cable whose weight equals the horizontal tension, the solution of the problem leads to the transcendental equation $s = \left(e^{\frac{l}{2z}} - e^{-\frac{l}{2z}} \right) z$, or inserting the numerical values,

$$f(z) = 22 - \left(e^{\frac{10}{z}} - e^{-\frac{10}{z}} \right) z = 0$$

is the equation to be solved. The first derivative equation is

$$f'(z) = - \left(e^{\frac{10}{z}} - e^{-\frac{10}{z}} \right) + \frac{10}{z} \left(e^{\frac{10}{z}} + e^{-\frac{10}{z}} \right) = 0,$$

and this substituted in $f(z)$ shows that one real root is less than about 20. Assume $z_1 = 15$, then $f(z_1) = 0.486$ and $f'(z_1) = 0.206$, whence by Newton's rule (Art. 4) $z_2 = 13$ nearly. Next for $z_2 = 13$, $f(z_2) = -0.0298$ and $f'(z_2) = 0.322$, whence $z_3 = 13.1$. Lastly for $z_3 = 13.1$ $f(z_3) = 0.0012$ and $f'(z_3) = 0.3142$, whence $z_4 = 13.096$, which is a sufficiently close approximation. The horizontal tension in the given catenary is hence 130.96 pounds.*

* Since $e^\theta - e^{-\theta} = 2 \sinh \theta$, this equation may be written $11\theta - 10 \sinh \theta$, where $\theta = 10z^{-1}$, and the solution may be expedited by the help of tables of hyperbolic functions. See Chapter IV.

Prob. 9. Show that the equation $3 \sin x - 2x - 5 = 0$ has but one real root, and compute its value.

Prob. 10. Find the number of real roots of the equation $2x + \log x - 10000 = 0$, and show that the value of one of them is $x = 4995.74$.

ART. 8. ALGEBRAIC SOLUTIONS.

Algebraic solutions of complete algebraic equations are only possible when the degree n is less than 5. It frequently happens, moreover, that the algebraic solution cannot be used to determine numerical values of the roots as the formulas expressing them are in irreducible imaginary form. Nevertheless the algebraic solutions of quadratic, cubic, and quartic equations are of great practical value, and the theory of the subject is of the highest importance, having given rise in fact to a large part of modern algebra.

The solution of the quadratic has been known from very early times, and solutions of the cubic and quartic equations were effected in the sixteenth century. A complete investigation of the fundamental principles of these solutions was, however, first given by Lagrange in 1770.* This discussion showed, if the general equation of the n^{th} degree, $f(x) = 0$, be deprived of its second term, thus giving the equation $f(y) = 0$, that the expression for the root y is given by

$$y = \omega s_1 + \omega^2 s_2 + \dots + \omega^{n-1} s_{n-1},$$

in which n is the degree of the given equation, ω is, in succession, each of the n^{th} roots of unity, $1, \epsilon, \epsilon^2, \dots, \epsilon^{n-1}$, and s_1, s_2, \dots, s_{n-1} , are the so-called elements which in soluble cases are determined by an equation of the $n - 1^{\text{th}}$ degree. For instance, if $n = 3$ the equation is of the third degree or a cubic, the three values of ω are

$$\omega_1 = 1, \quad \omega = -\frac{1}{2} + \frac{1}{2}\sqrt{-3} = \epsilon, \quad \omega = -\frac{1}{2} - \frac{1}{2}\sqrt{-3} = \epsilon^2,$$

* *Memoirs of Berlin Academy*, 1769 and 1770; reprinted in *Œuvres de Lagrange* (Paris, 1868), Vol. II, pp. 539-562. See also *Traité de la résolution des équations numériques*, Paris, 1798 and 1808.

and the three roots are expressed by

$$y_1 = s_1 + s_2, \quad y_2 = \epsilon s_1 + \epsilon^2 s_2, \quad y_3 = \epsilon^2 s_1 + \epsilon s_2,$$

in which s_1^3 and s_2^3 are found to be the roots of a quadratic equation (Art. 9).

The n values of ω are the n roots of the binomial equation $\omega^n - 1 = 0$. If n be odd, one of these is real and the others are imaginary; if n be even, two are real and $n - 2$ are imaginary.* Thus the roots of $\omega^3 - 1 = 0$ are $+1$ and -1 ; those of $\omega^5 - 1 = 0$ are given above; those of $\omega^4 - 1 = 0$ are $+1$, $+i$, -1 , and $-i$ where i is $\sqrt{-1}$. For the equation $\omega^5 - 1 = 0$ the real root is $+1$, and the imaginary roots are denoted by ϵ , ϵ^2 , ϵ^3 , ϵ^4 ; to find these let $\omega^5 - 1 = 0$ be divided by $\omega - 1$, giving

$$\omega^4 + \omega^3 + \omega^2 + \omega + 1 = 0,$$

which being a reciprocal equation can be reduced to a quadratic, and the solution of this furnishes the four values,

$$\epsilon = -\frac{1}{4}(1 - \sqrt{5} + \sqrt{-10 - 2\sqrt{5}}), \quad \epsilon^2 = -\frac{1}{4}(1 + \sqrt{5} + \sqrt{-10 + 2\sqrt{5}}),$$

$$\epsilon^4 = -\frac{1}{4}(1 - \sqrt{5} - \sqrt{-10 - 2\sqrt{5}}), \quad \epsilon^3 = -\frac{1}{4}(1 + \sqrt{5} - \sqrt{-10 + 2\sqrt{5}}),$$

where it will be seen that $\epsilon \cdot \epsilon^4 = 1$ and $\epsilon^2 \cdot \epsilon^3 = 1$, as should be the case, since $\epsilon^5 = 1$. *

In order to solve a quadratic equation by this general method let it be of the form

$$x^2 + 2ax + b = 0,$$

and let x be replaced by $y - a$, thus reducing it to

$$y^2 - (a^2 - b) = 0.$$

Now the two roots of this are $y_1 = +s_1$ and $y_2 = -s_1$, whence the product of $(y - s_1)$ and $(y + s_1)$ is

$$y^2 - s^2 = 0.$$

Thus the value of s^2 is given by an equation of the first degree,

* The values of ω are, in short, those of the n "vectors" drawn from the center which divide a circle of radius unity into n equal parts, the first vector $\omega_1 = 1$ being measured on the axis of real quantities. See Chapter X.

$s^2 = a^2 - b$; and since $x = -a + y$, the roots of the given equation are

$$x_1 = -a + \sqrt{a^2 - b}, \quad x_2 = -a - \sqrt{a^2 - b},$$

which is the algebraic solution of the quadratic.

The equation of the $n - 1^{\text{th}}$ degree upon which the solution of the equation of the n^{th} degree depends is called a resolvent. If such a resolvent exists, the given equation is algebraically solvable; but, as before remarked, this is only the case for quadratic, cubic, and quartic equations.

Prob. 11. Show that the six 6^{th} roots of unity are $+1$, $+\frac{1}{2}(1 + \sqrt{-3})$, $-\frac{1}{2}(1 - \sqrt{-3})$, -1 , $-\frac{1}{2}(1 + \sqrt{-3})$, $-\frac{1}{2}(1 - \sqrt{-3})$.

ART. 9. THE CUBIC EQUATION.

All methods for the solution of the cubic equation lead to the result commonly known as Cardan's formula.* Let the cubic be

$$x^3 + 3ax^2 + 3bx + 2c = 0, \quad (1)$$

and let the second term be removed by substituting $y - a$ for x , giving the form,

$$y^3 + 3By + 2C = 0, \quad (1')$$

in which the values of B and C are

$$B = -a^2 + b, \quad C = a^3 - \frac{3}{2}ab + c. \quad (2)$$

Now by the Lagrangian method of Art. 8 the values of y are

$$y_1 = s_1 + s_2, \quad y_2 = \epsilon s_1 + \epsilon^2 s_2, \quad y_3 = \epsilon^2 s_1 + \epsilon s_2,$$

in which ϵ and ϵ^2 are the imaginary cube roots of unity. Forming the products of the roots, and remembering that $\epsilon^3 = 1$ and $\epsilon^2 + \epsilon + 1 = 0$, there are found

$$\begin{aligned} y_1 y_2 + y_1 y_3 + y_2 y_3 &= -3s_1 s_2 = +3B, \\ y_1 y_2 y_3 &= s_1^3 + s_2^3 = -2C. \end{aligned}$$

For the determination of s_1 and s_2 there are hence two equations from which results the quadratic resolvent

$s^6 + 2Cs^3 - B^3 = 0$, and thus

$$s_1 = (-C + \sqrt{B^3 + C^2})^{\frac{1}{3}}, \quad s_2 = (-C - \sqrt{B^3 + C^2})^{\frac{1}{3}}. \quad (3)$$

* Deduced by Ferreo in 1515, and first published by Cardan in 1545.

One of the roots of the cubic in y therefore is

$$y_1 = (-C + \sqrt{B^3 + C^2})^{\frac{1}{3}} + (-C - \sqrt{B^3 + C^2})^{\frac{1}{3}},$$

and this is the well-known formula of Cardan.

The algebraic solution of the cubic equation (1) hence consists in finding B and C by (2) in terms of the given coefficients, and then by (3) the elements s_1 and s_2 are determined. Finally,

$$\begin{aligned} x_1 &= -a + (s_1 + s_2), \\ x_2 &= -a - \frac{1}{2}(s_1 + s_2) + \frac{1}{2}\sqrt{-3}(s_1 - s_2), \\ x_3 &= -a - \frac{1}{2}(s_1 + s_2) - \frac{1}{2}\sqrt{-3}(s_1 - s_2), \end{aligned} \quad (4)$$

which are the algebraic expressions of the three roots.

When $B^3 + C^2$ is negative the numerical solution of the cubic is not possible by these formulas, as then both s_1 and s_2 are in irreducible imaginary form. This, as is well known, is the case of three real roots, $s_1 + s_2$ being a real, while $s_1 - s_2$ is a pure imaginary.* When $B^3 + C^2$ is 0 the elements s_1 and s_2 are equal, and there are two equal roots, $x_2 = x_3 = -a + C^{\frac{1}{3}}$, while the other root is $x_1 = -a - 2C^{\frac{1}{3}}$.

When $B^3 + C^2$ is positive the equation has one real and two imaginary roots, and formulas (2), (3), and (4) furnish the numerical values of the roots of (1). For example, take the cubic

$$x^3 - 4.5x^2 + 12x - 5 = 0,$$

whence by comparison with (1) are found $a = -1.5$, $b = +4$, $c = -2.5$. Then from (2) are computed $B = 1.75$, $C = +3.125$. These values inserted in (3) give $s_1 = +0.9142$, $s_2 = -1.9142$; thus $s_1 + s_2 = -1.0$ and $s_1 - s_2 = +2.8284$. Finally, from (4)

$$\begin{aligned} x_1 &= 1.5 - 1.0 = +0.5, \\ x_2 &= 1.5 + 0.5 + 1.4142\sqrt{-3} = 2 + 2.4495i, \\ x_3 &= 1.5 + 0.5 - 1.4142\sqrt{-3} = 2 - 2.4495i, \end{aligned}$$

which are the three roots of the given cubic.

* The numerical solution of this case is possible whenever the angle whose cosine is $-C/\sqrt{-B^3}$ can be geometrically trisected.

Prob. 12. Compute the roots of $x^3 - 2x - 5 = 0$. Also the roots of $x^3 + 0.6x^2 - 5.76x + 4.32 = 0$.

Prob. 13. A cone has its altitude 6 inches and the diameter of its base 5 inches. It is placed with vertex downwards and one fifth of its volume is filled with water. If a sphere 4 inches in diameter be then put into the cone, what part of its radius is immersed in the water? (Ans. 0.5459 inches).

ART. 10. THE QUARTIC EQUATION.

The quartic equation was first solved in 1545 by Ferrari, who separated it into the difference of two squares. Descartes in 1637 resolved it into the product of two quadratic factors. Tschirnhausen in 1683 removed the second and fourth terms. Euler in 1732 and Lagrange in 1767 effected solutions by assuming the form of the roots. All these methods lead to cubic resolvents, the roots of which are first to be found in order to determine those of the quartic.

The methods of Euler and Lagrange, which are closely similar, first reduce the quartic to one lacking the second term,

$$y^4 + 6By^2 + 4Cy + D = 0;$$

and the general form of the roots being taken as

$$\begin{aligned} y_1 &= +\sqrt{s_1} + \sqrt{s_2} + \sqrt{s_3}, & y_3 &= -\sqrt{s_1} + \sqrt{s_2} - \sqrt{s_3}, \\ y_2 &= +\sqrt{s_1} - \sqrt{s_2} - \sqrt{s_3}, & y_4 &= -\sqrt{s_1} - \sqrt{s_2} + \sqrt{s_3}, \end{aligned}$$

the values s_1, s_2, s_3 , are shown to be the roots of the resolvent,

$$s^3 + 3Bs^2 + \frac{1}{4}(9B^2 - D)s - \frac{1}{4}C^2 = 0.$$

Thus the roots of the quartic are algebraically expressed in terms of the coefficients of the quartic, since the resolvent is solvable by the process of Art. 9.

Whatever method of solution be followed, the following final formulas, deduced by the author in 1892, will result.* Let the complete quartic equation be written in the form

$$x^4 + 4ax^3 + 6bx^2 + 4cx + d = 0. \quad (1)$$

* See American Journal Mathematics, 1892, Vol. XIV, pp. 237-245.

First, let g , h , and k be determined from

$$g = a^3 - b, \quad h = b^3 + c^2 - 2abc + dg, \quad k = \frac{4}{3}ac - b^3 - \frac{1}{3}d. \quad (2)$$

Secondly, let l be obtained by

$$l = \frac{1}{2}(h + \sqrt{h^2 + k^3})^{\frac{1}{3}} + \frac{1}{2}(h - \sqrt{h^2 + k^3})^{\frac{1}{3}} \quad (3)$$

Thirdly, let u , v , and w be found from

$$u = g + l, \quad v = 2g - l, \quad w = 4u^3 + 3k - 12gl. \quad (4)$$

Then the four roots of the quartic equation are

$$\left. \begin{aligned} x_1 &= -a + \sqrt{u} + \sqrt{v + \sqrt{w}}, \\ x_2 &= -a + \sqrt{u} - \sqrt{v + \sqrt{w}}, \\ x_3 &= -a - \sqrt{u} + \sqrt{v - \sqrt{w}}, \\ x_4 &= -a - \sqrt{u} - \sqrt{v - \sqrt{w}}, \end{aligned} \right\} \quad (5)$$

in which the signs are to be used as written provided that $2a^3 - 3ab + c$ is a negative number; but if this is positive all radicals except \sqrt{w} are to be reversed in sign.

These formulas not only serve for the complete theoretic discussion of the quartic (1), but they enable numerical solutions to be made whenever (3) can be computed, that is, whenever $h^2 + k^3$ is positive. For this case the quartic has two real and two imaginary roots. If there be either four real roots or four imaginary roots $h^2 + k^3$ is negative, and the irreducible case arises where convenient numerical values cannot be obtained, although they are correctly represented by the formulas.

As an example let a given rectangle have the sides p and q , and let it be required to find the length of an inscribed rectangle whose width is m . If x be this length, this is a root of the quartic equation

$$x^4 - (p^2 + q^2 + 2m^2)x^2 + 4pqmx - (p^2 + q^2 - m^2)m^2 = 0,$$

and thus the problem is numerically solvable by the above formulas if two roots are real and two imaginary. As a special case let $p = 4$ feet, $q = 3$ feet, and $m = 1$ foot; then

$$x^4 - 27x^2 + 48x - 24 = 0.$$

By comparison with (1) are found $a = 0$, $b = -4\frac{1}{2}$, $c = +12$, and $d = -24$. Then from (2), $g = +4\frac{1}{2}$, $h = -4\frac{1}{8}$, and $k = +\frac{1}{4}$. Thus $h^2 + k^3$ is positive, and from (3) the value of l is -3.6067 . From (4) are now found, $u = +0.8933$, $v = 12.6067$, and $w = +161.20$. Then, since c is positive, the values of the four roots are, by (5),

$$\begin{aligned}x_1 &= -0.945 - \sqrt{12.607 + 12.697} = -5.975 \text{ feet,} \\x_2 &= -0.945 + \sqrt{12.607 + 12.697} = +4.085 \text{ feet,} \\x_3 &= +0.945 - \sqrt{12.607 - 12.697} = +0.945 - 0.30i, \\x_4 &= +0.945 + \sqrt{12.607 - 12.697} = +0.945 + 0.30i,\end{aligned}$$

the second of which is evidently the required length. Each of these roots closely satisfies the given equation, the slight discrepancy in each case being due to the rounding off at the third decimal.*

Prob. 14. Compute the roots of the equation $x^4 + 7x + 6 = 0$. (Ans. -1.388 , -1.000 , $1.194 \pm 1.701i$.)

ART. 11. QUINTIC EQUATIONS.

The complete equation of the fifth degree is not algebraically solvable, nor is it reducible to a solvable form. Let the equation be

$$x^5 + 5ax^4 + 5bx^3 + 5cx^2 + 5dx + 2e = 0,$$

and by substituting $y - a$ for x let it be reduced to

$$y^5 + 5By^3 + 5Cy^2 + 5Dy + 2E = 0.$$

The five roots of this are, according to Art. 8,

$$\begin{aligned}y_1 &= s_1 + s_2 + s_3 + s_4, \\y_2 &= \epsilon s_1 + \epsilon^2 s_2 + \epsilon^3 s_3 + \epsilon^4 s_4, \\y_3 &= \epsilon^2 s_1 + \epsilon^4 s_2 + \epsilon s_3 + \epsilon^3 s_4, \\y_4 &= \epsilon^3 s_1 + \epsilon s_2 + \epsilon^4 s_3 + \epsilon^2 s_4, \\y_5 &= \epsilon^4 s_1 + \epsilon^3 s_2 + \epsilon^2 s_3 + \epsilon s_4,\end{aligned}$$

in which ϵ , ϵ^2 , ϵ^3 , ϵ^4 are the imaginary fifth roots of unity. Now if the several products of these roots be taken there will be

* This example is known by civil engineers as the problem of finding the length of a strut in a panel of the Howe truss.

found, by (4) of Art. 6, four equations connecting the four elements s_1, s_2, s_3 , and s_4 , namely,

$$- B = s_1 s_4 + s_2 s_3,$$

$$- C = s_1^2 s_3 + s_2^2 s_1 + s_3^2 s_4 + s_4^2 s_2,$$

$$- D = s_1^3 s_2 + s_2^3 s_4 + s_3^3 s_1 + s_4^3 s_3 - s_1^2 s_4^2 - s_2^2 s_3^2 + s_1 s_2 s_3 s_4,$$

$$- 2E = s_1^5 + s_2^5 + s_3^5 + s_4^5 + 5(s_1^2 s_2^2 s_4 + s_1^2 s_3^2 s_2 + s_2^2 s_4^2 s_3 + s_3^2 s_4^2 s_1) \\ - 5(s_1^3 s_3 s_4 + s_2^3 s_1 s_3 + s_3^3 s_2 s_4 + s_4^3 s_1 s_2);$$

but the solution of these leads to an equation of the 120th degree for s , or of the 24th degree for s^5 . However, by taking $s_1 s_4 - s_2 s_3$ or $s_1^5 + s_2^5 + s_3^5 + s_4^5$ as the unknown quantity, a resolvent of the 6th degree is obtained, and all efforts to find a resolvent of the fourth degree have proved unavailing.

Another line of attack upon the quintic is in attempting to remove all the terms intermediate between the first and the last. By substituting $y^2 + py + q$ for x , the values of p and q may be determined so as to remove the second and third terms by a quadratic equation, or the second and third by a cubic equation, or the second and fourth by a quartic equation, as was first shown by Tschirnhausen in 1683. By substituting $y^3 + py^2 + qy + r$ for x , three terms may be removed, as was shown by Bring in 1786. By substituting $y^4 + py^3 + qy^2 + ry + t$ for x it was thought by Jerrard in 1833 that four terms might be removed, but Hamilton showed later that this leads to equations of a degree higher than the fourth.

In 1826 Abel gave a demonstration that the algebraic solution of the general quintic is impossible, and later Galois published a more extended investigation leading to the same conclusion.* The reason for the algebraic solvability of the quartic equation may be briefly stated as the fact that there exist rational three-valued functions of four quantities. There are, however, no rational four-valued functions of five quantities, and accordingly a quartic resolvent cannot be found for the general quintic equation.

* Jordan's *Traité des substitutions et des équations algébriques*; Paris, 1870. *Abhandlungen über die algebraische Auflösung der Gleichungen* von N. H. Abel und Galois; Berlin, 1889.

There are, however, numerous special forms of the quintic whose algebraic solution is possible. The oldest of these is the quintic of De Moivre,

$$y^5 + 5By^3 + 5B^2y + 2E = 0,$$

which is solved at once by making $s_2 = s_3 = 0$ in the element equations; then $-B = s_1s_4$ and $-2E = s_1^5 + s_4^5$, from which s_1 and s_4 are found, and $y_1 = s_1 + s_4$, or

$$y_1 = (-E + \sqrt{B^5 + E^2})^{\frac{1}{5}} + (-E - \sqrt{B^5 + E^2})^{\frac{1}{5}},$$

while the other roots are $y_2 = \epsilon s_1 + \epsilon^4 s_4$, $y_3 = \epsilon^2 s_1 + \epsilon^3 s_4$, $y_4 = \epsilon^3 s_1 + \epsilon^2 s_4$, and $y_5 = \epsilon^4 s_1 + \epsilon s_4$. If $B^5 + E^2$ be negative, this quintic has five real roots; if positive, there are one real and four imaginary roots.

When any relation, other than those expressed by the four element equations, exists between s_1, s_2, s_3, s_4 , the quintic is solvable algebraically. As an infinite number of such relations may be stated, it follows that there are an infinite number of solvable quintics. In each case of this kind, however, the coefficients of the quintic are also related to each other by a certain equation of condition.

The complete solution of the quintic in terms of one of the roots of its resolvent sextic was made by McClintock in 1884.* By this method s_1^5, s_2^5, s_3^5 , and s_4^5 are expressed as the roots of a quartic in terms of a quantity t which is the root of a sextic whose coefficients are rational functions of those of the given quintic. Although this has great theoretic interest, it is, of course, of little practical value for the determination of numerical values of the roots.

By means of elliptic functions the complete quintic can, however, be solved, as was first shown by Hermite in 1858. For this purpose the quintic is reduced by Jerrard's transformation to the form $x^5 + 5dx + 2e = 0$, and to this form can also be reduced the elliptic modular equation of the sixth degree. Other solutions by elliptic functions were made by

* American Journal of Mathematics, 1886, Vol. VIII, pp. 49-83.

Kronecker in 1861 and by Klein in 1884.* These methods, though feasible by the help of tables, have not yet been systematized so as to be of practical advantage in the numerical computation of roots.

Prob. 15. If the relation $s_1 s_4 = s_2 s_3$ exists between the elements show that $s_1^6 + s_2^6 + s_3^6 + s_4^6 = -2E$.

Prob. 16. Compute the roots of $y^5 + 10y^3 + 20y + 6 = 0$, and also those of $y^5 - 10y^3 + 20y + 6 = 0$.

ART. 12. TRIGONOMETRIC SOLUTIONS.

When a cubic equation has three real roots the most convenient practical method of solution is by the use of a table of sines and cosines. If the cubic be stated in the form (1) of Art. 9, let the second term be removed, giving

$$y^3 + 3By + 2C = 0.$$

Now suppose $y = 2r \sin \theta$, then this equation becomes

$$8 \sin^3 \theta + 6 \frac{B}{r^2} \sin \theta + 2 \frac{C}{r^3} = 0,$$

and by comparison with the known trigonometric formula

$$8 \sin^3 \theta - 6 \sin \theta + 2 \sin 3\theta = 0,$$

there are found for r and $\sin 3\theta$ the values

$$r = \sqrt{-B}, \quad \sin 3\theta = C / \sqrt{-B^3},$$

in which B is always negative for the case of three real roots (Art. 9). Now $\sin 3\theta$ being computed, 3θ is found from a table of sines, and then θ is known. Thus,

$y_1 = 2r \sin \theta$, $y_2 = 2r \sin (120^\circ + \theta)$, $y_3 = 2r \sin (240^\circ + \theta)$, are the real roots of the cubic in y .†

* For an outline of these transcendental methods, see Hagen's *Synopsis der höheren Mathematik*, Vol. I, pp. 339-344.

† When B^3 is negative and numerically less than C^2 , as also when B^3 is positive, this solution fails, as then one root is real and two are imaginary. In this case, however, a similar method of solution by means of hyperbolic sines is possible. See Grunert's *Archiv für Mathematik und Physik*, Vol. xxxviii, pp. 48-76.

For example, the depth of flotation of a sphere whose diameter is 2 feet and specific gravity 0.65, is given by the cubic equation $x^3 - 3x^2 + 2.6 = 0$ (Art. 6). Placing $x = y + 1$ this reduces to $y^3 - 3y + 0.6 = 0$, for which $B = -1$ and $C = +0.3$. Thus $r = 1$ and $\sin 3\theta = +0.3$. Next from a table of sines, $3\theta = 17^\circ 27'$, and accordingly $\theta = 5^\circ 49'$. Then

$$y_1 = 2 \sin 5^\circ 49' = +0.2027,$$

$$y_2 = 2 \sin 125^\circ 49' = +1.6218,$$

$$y_3 = 2 \sin 245^\circ 49' = -1.8245.$$

Adding 1 to each of these, the values of x are

$$x_1 = +1.203 \text{ feet}, \quad x_2 = +2.622 \text{ feet}, \quad x_3 = -0.825 \text{ feet};$$

and evidently, from the physical aspect of the question, the first of these is the required depth. It may be noted that the number 0.3 is also the sine of $162^\circ 11'$, but by using this the three roots have the same values in a different order.

When the quartic equation has four real roots its cubic resolvent has also three real roots. In this case the formulas of Art. 10 will furnish the solution if the three values of l be obtained from (3) by the help of a table of sines. The quartic being given, g , h , and k are found as before, and the value of k will always be negative for four real roots. Then

$$r = \sqrt{-k}, \quad \sin 3\theta = -h/r^3,$$

and 3θ is taken from a table; thus θ is known, and the three values of l are

$$l_1 = r \sin \theta, \quad l_2 = r \sin(120^\circ + \theta), \quad l_3 = r \sin(240^\circ + \theta).$$

Next the three values of u , of v , and of w are computed, and those selected which give u , w , and $v - \sqrt{w}$ all positive quantities. Then (5) gives the required roots of the quartic.

As an example, take the case of the inscribed rectangle in Art. 10, and let $p = 4$ feet, $q = 3$ feet, $m = \sqrt{13}$ feet; then the quartic equation is

$$x^4 - 51x^2 + 48\sqrt{13}x - 156 = 0.$$

Here $a = 0$, $b = -8\frac{1}{2}$, $c = +12\sqrt{13}$, and $d = -156$. Next $g = +8\frac{1}{2}$, $h = -\frac{545}{8}$, and $k = -\frac{81}{4}$. The trigonometric work now begins; the value of r is found to be $+4\frac{1}{2}$, and that of $\sin 3\theta$ to be $+0.7476$; hence from the table $3\theta = 48^\circ 23'$, and $\theta = 16^\circ 07' 40''$. The three values of l are then computed by logarithmic tables, and found to be,

$$l_1 = +1.250, \quad l_2 = +3.1187, \quad l_3 = -4.3687.$$

Next the values of u , v , and w are obtained, and it is seen that only those corresponding to l_1 will render all quantities under the radicals positive; these quantities are $u = 9.75$, $v = 15.75$, and $w = 192.0$. Then the four roots of the quartic are

$x_1 = -8.564$, $x_2 = +2.319$, $x_3 = +1.746$, $x_4 = +4.499$ feet, of which only the second and third belong to inscribed rectangles, while the first and fourth belong to rectangles whose corners are on the sides of the given rectangle produced.

Trigonometric solutions of the quintic equation are not possible except for the binomial $x^5 \pm a$, and the quintic of De Moivre. The general trigonometric expression for the root of a quintic lacking its second term is $y = 2r_1 \cos \theta_1 + 2r_2 \cos \theta_2$, and to render a solution possible, r_1 and r_2 , as well as $\cos \theta_1$ and $\cos \theta_2$, must be found; but these in general are roots of equations of the sixth or twelfth degree: in fact r_1^2 is the same as the function $s_1 s_4$ of Art. 11, and r_2^2 is the same as $s_2 s_3$. Here $\cos \theta_1$ and $\cos \theta_2$ may be either circular or hyperbolic cosines, depending upon the signs and values of the coefficients of the quintic.

Trigonometric solutions are possible for any binomial equation, and also for any equation which expresses the division of an angle into equal parts. Thus the roots of $x^5 + 1 = 0$ are $\cos m 30^\circ \pm i \sin m 30^\circ$, in which m has the values 1, 2, and 3. The roots of $x^5 - 5x^3 + 5x - 2 \cos 5\theta = 0$ are $2 \cos (m 72^\circ + \theta)$ where m has the values 0, 1, 2, 3, and 4.

Prob. 17. Compute by a trigonometric solution the four roots of the quartic $x^4 + 4x^3 - 24x^2 - 76x - 29 = 0$. (Ans. -6.734 , -1.550 , $+0.262$, $+4.022$).

Prob. 18. Give a trigonometric solution of the quintic equation $x^5 - 5bx^3 + 5b^2x - 2e = 0$ for the case of five real roots. Compute the roots when $b = 1$ and $e = 0.752798$. (Ans. -1.7940 , -1.3952 , 0.2864 , 0.9317 , 1.9710 .)

ART. 13. REAL ROOTS BY SERIES.

The value of x in any algebraic equation may be expressed as an infinite series. Let the equation be of any degree, and by dividing by the coefficient of the term containing the first power of x let it be placed in the form

$$a = x + bx^2 + cx^3 + dx^4 + ex^5 + fx^6 + \dots$$

Now let it be assumed that x can be expressed by the series

$$x = a + ma^2 + na^3 + pa^4 + qa^5 + \dots$$

By inserting this value of x in the equation and equating the coefficients of like powers of a , the values of m , n , etc., are found, and then

$$x = a - ba^2 + (2b^2 - c)a^3 - (5b^3 - 5bc + d)a^4 + (14b^4 - 21b^2c + 6bd + 3c^2 - e)a^5 - (42b^5 - 84b^3c + 28b^2d + 28bc^2 - 7be - 7cd + f)a^6 + \dots,$$

is an expression of one of the roots of the equation. In order that this series may converge rapidly it is necessary that a should be a small fraction.*

To apply this to a cubic equation the coefficients d , e , f , etc., are made equal to 0. For example, let $x^3 - 3x + 0.6 = 0$; this reduced to the given form is $0.2 = x - \frac{1}{3}x^3$, hence $a = 0.2$, $b = 0$, $c = -\frac{1}{3}$, and then

$$x = 0.2 + \frac{1}{3} \cdot 0.2^3 + \frac{1}{3} \cdot 0.2^5 + \text{etc.} = +0.20277,$$

which is the value of one of the roots correct to the fourth decimal place. This equation has three real roots, but the series gives only one of them; the others can, however, be found if their approximate values are known. Thus, one root is about $+1.6$, and by placing $x = y + 1.6$ there results an equation in y whose root by the series is found to be $+0.0218$, and hence $+1.6218$ is another root of $x^3 - 3x + 0.6 = 0$.

* This method is given by J. B. Mott in *The Analyst*, 1882, Vol. IX, p. 104.

Cardan's expression for the root of a cubic equation can be expressed as a series by developing each of the cube roots by the binomial formula and adding the results. Let the equation be $y^3 + 3By + 2C = 0$, whose root is, by Art. 9,

$$y = (-C + \sqrt{B^3 + C^2})^{\frac{1}{3}} + (-C - \sqrt{B^3 + C^2})^{\frac{1}{3}},$$

then this development gives the series,

$$y = 2(-C)^{\frac{1}{3}} \left(1 - \frac{2}{2}r - \frac{2 \cdot 5 \cdot 8}{2 \cdot 3 \cdot 4}r^2 - \frac{2 \cdot 5 \cdot 8 \cdot 11 \cdot 14}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6}r^3 - \dots \right),$$

in which r represents the quantity $(B^3 + C^2)/3C^2$. If $r = 0$ the equation has two equal roots and the third root is $2(-C)^{\frac{1}{3}}$. If r is numerically greater than unity the series is divergent, and the solution fails. If r is numerically less than unity and sufficiently small to make a quick convergence, the series will serve for the computation of one real root. For example, take the equation $x^3 - 6x + 6 = 0$, where $B = -2$ and $C = 3$; hence $r = 1/81$, and one root is

$y = -2.8845(1 - 0.01235 - 0.00051 - 0.00032 -) = -2.846$, which is correct to the third decimal. In comparatively few cases, however, is this series of value for the solution of cubics.

Many other series for the expression of the roots of equations, particularly for trinomial equations, have been devised. One of the oldest is that given by Lambert in 1758, whereby the root of $x^n + ax - b = 0$ is developed in terms of the ascending powers of b/a . Other solutions were published by Euler and Lagrange. These series usually give but one root, and this only when the values of the coefficients are such as to render convergence rapid.

Prob. 19. Consult Euler's *Anleitung zur Algebra* (St. Petersburg, 1771), pp. 143-150, and apply his method of series to the solution of a quartic equation.

ART. 14. COMPUTATION OF ALL ROOTS.

A comprehensive and valuable method for the solution of equations by series was developed by McClintock, in 1894, by

means of his Calculus of Enlargement.* By this method all the roots, whether real or imaginary, may be computed from a single series. The following is a statement of the method as applied to trinomial equations:

Let $x^n = nAx^{n-k} + B^n$ be the given trinomial equation. Substitute $x = By$ and thus reduce the equation to the form $y^n = nay^{n-k} + 1$ where $a = A/B^k$. Then if B^n is positive, the roots are given by the series

$$\begin{aligned} y = \omega + \omega^{1-k}a + \omega^{1-2k}(1 - 2k + n)a^2/2! \\ + \omega^{1-3k}(1 - 3k + n)(1 - 3k + 2n)a^3/3! \\ + \omega^{1-4k}(1 - 4k + n)(1 - 4k + 2n)(1 - 4k + 3n)a^4/4! + \dots, \end{aligned}$$

in which ω represents in succession each of the roots of unity. If, however, B^n is negative, the given equation reduces to $y^n = nay^{n-k} - 1$, and the same series gives the roots if ω be taken in succession as each of the roots of -1 .

In order that this series may be convergent the value of a^n must be numerically less than $k^{-k}(n-k)^{k-n}$; thus for the quartic $y^4 = 4ax + 1$, where $n = 4$ and $k = 3$, the value of a must be less than $27^{-\frac{1}{4}}$.

To apply this method to the cubic equation $x^3 = 3Ax \pm B^3$, place $n = 3$ and $k = 2$, and put $y = Bx$. It then becomes $y^3 = 3ay \pm 1$ where $a = A/B^2$, and the series is

$$y = \omega + \omega^2a - \frac{1}{3}\omega a^3 + \frac{1}{3}\omega^2a^4 + \dots,$$

in which the values to be taken for ω are the cube roots of 1 or -1 , as the case may be. For example, let $x^3 - 2x - 5 = 0$. Placing $y = 5^{\frac{1}{3}}x$, this reduces to $y^3 = 0.684y + 1$. Here $a = 0.228$, and as this is less than $4^{-\frac{1}{3}}$ the series is convergent. Making $\omega = 1$, the first root is

$$y = 1 + 0.2280 - 0.0039 + 0.0009 = 1.2250.$$

* See Bulletin of American Mathematical Society, 1894, Vol. I, p. 3; also American Journal of Mathematics, 1895, Vol. XVII, pp. 89-110.

Next making $\omega = -\frac{1}{2} + \frac{1}{2}\sqrt{-3}$, ω^2 is $-\frac{1}{2} - \frac{1}{2}\sqrt{-3}$, and the corresponding root is found to be

$$y = -0.6125 + 0.3836\sqrt{-3}.$$

Again, making $\omega = -\frac{1}{2} - \frac{1}{2}\sqrt{-3}$ the third root is found to be the conjugate imaginary of the second. Lastly, multiplying each value of y by $5^{\frac{1}{3}}$,

$$x = 2.095, \quad x = -1.047 \pm 1.136\sqrt{-1},$$

which are very nearly the roots of $x^3 - 2x - 5 = 0$.

In a similar manner the cubic $x^3 + 2x + 5 = 0$ reduces to $y^3 = -0.684y - 1$, for which the series is convergent. Here the three values of ω are, in succession, -1 , $\frac{1}{2} + \frac{1}{2}\sqrt{-3}$, $-\frac{1}{2} + \frac{1}{2}\sqrt{-3}$, and the three roots are $y = -0.777$ and $y = 0.388 \pm 1.137i$.

When all the roots are real, the method as above stated fails because the series is divergent. The given equation can, however, be transformed so as to obtain $n - k$ roots by one application of the general series and k roots by another. As an example, let $x^3 - 243x + 330 = 0$. For the first application this is to be written in the form

$$x = \frac{x^3}{243} + \frac{330}{243},$$

for which $n = 1$ and $k = -2$. To make the last term unity place $x = \frac{330}{243}y$, and the equation becomes

$$y = \frac{330^3}{243^3}y^3 + 1,$$

whence $a = 330^3/3.243^3$. These values of n , k , and a are now inserted in the above general value of y , and ω made unity; thus $y = 0.9983$, whence $x_1 = 1.368$ is one of the roots. For the second application the equation is to be written

$$x^3 = -\frac{330}{243}x^{-1} + 243,$$

for which $n = 2$ and $k = 3$. Placing $x = 243^{\frac{1}{3}}y$, this becomes

$$y^3 = -\frac{340}{243^{\frac{1}{3}}}y^{-1} + 1,$$

whence $a = -110/243^{\frac{1}{3}}$, and the series is convergent. These values of n , k , and a are now inserted in the formula for y , and ω is made $+1$ and -1 in succession, thus giving two values for y , from which $x_2 = 14.86$ and $x_3 = -16.22$ are the other roots of the given cubic.

McClintock has also given a similar and more general method applicable to other algebraic equations than trinomials. The equation is reduced to the form $y^n = na \cdot \phi y \pm 1$, where $na \cdot \phi y$ denotes all the terms except the first and the last. Then the values of y are expressed by the series

$$y = \omega + \omega^{1-n} \phi \omega \cdot a + \omega^{1-n} \frac{d}{d\omega} \omega^{1-n} (\phi \omega)^2 \cdot \frac{a^2}{2!} + \\ + \left(\omega^{1-n} \frac{d}{d\omega} \right)^2 \omega^{1-n} (\phi \omega)^3 \cdot \frac{a^3}{3!} + \dots,$$

in which the values of ω are to be taken as before. The method is one of great importance in the theory of equations, as it enables not only the number of real and imaginary roots to be determined, but also gives their values when the convergence of the series is secured.

Prob. 20. Compute by the above method all the roots of the quartic $x^4 + x + 10 = 0$.

ART. 15. ROOTS OF UNITY.

The roots of $+1$ and -1 are required to be known in the numerical solution of algebraic equations by the method of the last article. From the theory of binomial equations given in all text-books on algebra, the n roots of $+1$ are

$$(+1)^{\frac{m}{n}} = \cos (m/n)2\pi + i \sin (m/n)2\pi, \quad m = 1, 2, 3, \dots n, \quad (1)$$

while those of -1 are expressed by

$$(-1)^{\frac{m}{n}} = \cos (m/n)\pi + i \sin (m/n)\pi, \quad m = 1, 2, 3, \dots n, \quad (2)$$

in which i represents the square root of -1 . From these general formulas it is seen that the two imaginary cube roots of $+1$ are

$$\epsilon_1 = -\frac{1}{2} + \frac{1}{2}i\sqrt{3} = -0.5 + 0.8660254i,$$

$$\epsilon_2 = -\frac{1}{2} - \frac{1}{2}i\sqrt{3} = -0.5 - 0.8660254i,$$

and that the two imaginary cube roots of -1 are

$$\epsilon_1' = +\frac{1}{2} + \frac{1}{2}i\sqrt{3} = +0.5 + 0.8660254i,$$

$$\epsilon_2' = +\frac{1}{2} - \frac{1}{2}i\sqrt{3} = +0.5 - 0.8660254i.$$

For the first case $\epsilon_1 + \epsilon_2 + 1 = 0$ and $\epsilon_1\epsilon_2 = 1$, as also $\epsilon_1 = \epsilon_2^2$ and $\epsilon_2^2 = \epsilon_1$, and similar relations apply to the other case.

The imaginary fifth roots of positive unity are given in Art. 8 expressed in radicals; reducing these to decimals, or deriving them from the above formula (1) with the help of a trigonometric table, there result

$$\epsilon = +0.3090170 + 0.9510565i, \quad \epsilon^2 = -0.8090170 + 0.5877853i,$$

$$\epsilon^4 = +0.3090170 - 0.9510565i, \quad \epsilon^3 = -0.8090170 - 0.5877853i,$$

while the imaginary fifth roots of negative unity are obtained from these by changing the signs. In general, if ω is an imaginary n^{th} root of positive unity, $-\omega$ is an imaginary n^{th} root of negative unity.

The imaginary sixth roots of positive unity may be expressed in terms of the cube roots. Let ϵ be one of the imaginary cube roots of $+1$, then the imaginary sixth roots of $+1$ are $+\epsilon$, $+\epsilon^2$, $-\epsilon$, $-\epsilon^2$; these are also the imaginary sixth roots of -1 .

From (1) the imaginary seventh roots of $+1$ are found to be

$$\epsilon = +0.6234898 + 0.7818316i, \quad \epsilon^6 = +0.6234898 - 0.7818316i,$$

$$\epsilon^2 = -0.2225209 + 0.9749234i, \quad \epsilon^5 = -0.2225209 - 0.9749234i,$$

$$\epsilon^3 = -0.9009688 + 0.4338837i, \quad \epsilon^4 = -0.9009688 - 0.4338837i,$$

and if the signs of these be reversed there result the imaginary seventh roots of -1 .

The imaginary eighth roots of $+1$ are $+i$, $-i$, $+\frac{1}{2}\sqrt{2}(1 \pm i)$, and $-\frac{1}{2}\sqrt{2}(1 \pm i)$. The imaginary ninth roots of $+1$ are the two

imaginary cube roots of $+1$, $\cos \frac{2}{3}\pi \pm i \sin \frac{2}{3}\pi$, and $\cos \frac{4}{3}\pi \pm i \sin \frac{4}{3}\pi$. The imaginary tenth roots of $+1$ are the five imaginary roots of $+1$ and the five imaginary roots of -1 . For any value of n the roots of $+1$ may be graphically represented in a circle of unit radius by taking one radius as $+1$ and drawing other radii to divide the circle into n equal parts; if unit distances normal to $+1$ and -1 be called $+i$ and $-i$, the n radii represent all the roots of $+1$. When this figure is viewed in a mirror, the image represents the n roots of -1 . Or, in other words, the $(m/n)^{\text{th}}$ roots of $+1$ are unit vectors which make the angles $(m/n)2\pi$ with the unit vector $+1$, while the $(m/n)^{\text{th}}$ roots of -1 are unit vectors which make the angles $(m/n)2\pi$ with the unit vector -1 .

The n roots of any unit vector $\cos \theta + i \sin \theta$ are readily found from De Moivre's theorem by the help of trigonometric tables. Accordingly the cube roots of this vector are $\cos \frac{1}{3}\theta + i \sin \frac{1}{3}\theta$, $\cos \frac{1}{3}(\theta + 2\pi) + i \sin \frac{1}{3}(\theta + 2\pi)$ and $\cos \frac{1}{3}(\theta + 4\pi) + i \sin \frac{1}{3}(\theta + 4\pi)$; the vectors representing these three roots divide the circle into three parts. The trigonometric solution of the cubic equation (Art. 12) is one application of De Moivre's theorem.

Prob. 21. Compute to six decimal places two or more of the eleventh imaginary roots of unity.

Prob. 22. Compute to five decimal places the five roots of the equation $x^5 - 0.8 - 0.6i = 0$.

Prob. 23. Compute to five decimal places the six roots of the equation $x^6 - 80 + 60i = 0$.

ART. 16. SOLUTIONS BY MACLAURIN'S FORMULA.

In 1903 Lambert published a method for the expression by Maclaurin's formula of the roots of equations in infinite series.* It applies to both algebraic and transcendental equations, and for the former it gives all the roots whether they be real or imaginary. The method is based on the device of introducing a

* Proceedings American Philosophical Society, Vol. 42.

factor x into all the terms but two of the equation $f(y)=0$, whereby y becomes an implicit function of x . The successive derivatives of y with respect to x are then obtained, and their values, as also those of y , are evaluated for $x=0$. By Maclaurin's formula, the expansions of y in powers of x become known, and if x be made unity in these expansions, the roots of $f(y)=0$ are found, provided the resulting series are convergent.

To illustrate this method by a numerical example, take the quartic equation

$$y^4 - 3y^2 + 75y - 10\,000 = 0, \quad (1)$$

and introduce an x into the second and third terms, thus,

$$y^4 - 3xy^2 + 75xy - 10\,000 = 0. \quad (2)$$

By Maclaurin's formula y may be expressed in terms of x , and then when x is made unity, the four series thus obtained furnish the four roots of (1). Maclaurin's formula is

$$y = y_0 + \left(\frac{dy}{dx}\right)_0 x + \left(\frac{d^2y}{dx^2}\right)_0 \frac{x^2}{2!} + \left(\frac{d^3y}{dx^3}\right)_0 \frac{x^3}{3!} + \dots,$$

where y_0 , $(dy/dx)_0$, $(d^2y/dx^2)_0$, etc., denote the values which y and the successive derivatives take when x is made 0. Differentiating equation (2) twice in succession, and then placing $x=0$, there are found

$$\begin{aligned} y_0 &= +10, & +10, & +10i, & -10i, \\ (dy/dx)_0 &= -0.1125, & -0.2625, & +0.1875 - 0.0750i, & +0.1875 + 0.0750i, \\ (d^2y/dx^2)_0 &= -0.0030, & +0.0030, & -0.0000 + 0.0039i, & -0.0000 - 0.0039i, \end{aligned}$$

in which i represents the square root of negative unity. Substituting each set of corresponding values in Maclaurin's formula and then placing $x=1$, there result

$$\begin{aligned} y_1 &= +9.886, & y_3 &= 0.1875 + 9.927i, \\ y_2 &= -10.261, & y_4 &= 0.1875 - 9.927i, \end{aligned}$$

which are the roots of (1), all correct to the last decimal.

This method may be readily applied to the trinomial equation $y^n - n a y^{n-k} - b = 0$. When x is inserted in the second term, the series obtained is

$$\begin{aligned} y = & b^{\frac{1}{n}} + \left(b^{\frac{1}{n}}\right)^{1-k} a + \left(b^{\frac{1}{n}}\right)^{1-2k} (1-2k+n) a^2 / 2! \\ & + \left(b^{\frac{1}{n}}\right)^{1-3k} (1-3k+n)(1-3k+2n) a^3 / 3! \\ & + \left(b^{\frac{1}{n}}\right)^{1-4k} (1-4k+n)(1-4k+2n)(1-4k+3n) a^4 / 4! + \dots \end{aligned}$$

and each of the roots is hence expressed in an infinite series, since $b^{\frac{1}{n}}$ has n values. This series is convergent when a^n is numerically less than $k^{-k}(n-k)^{k-n}b^k$, and for this case the roots can be computed. Now the condition $a^n = k^{-k}(n-k)^{k-n}b^k$ is that of equal roots in the trinomial equation; hence for the cubic equation the above series is applicable when one root is real and the others imaginary, while for the quartic equation it is applicable when two roots are real and two imaginary. For the irreducible case in cubics and quartics the above series does not converge and the roots cannot be computed from it; this case is treated on the next page by inserting x in other terms. This series is the same as that derived for trinomial equations by McClintock's method of enlargement (Art. 14).

As a special case take the quintic equation $y^5 - 5ay - 1 = 0$, in which the value of n is 5, that of k is 4, and those of $b^{\frac{1}{5}}$ are the five imaginary roots of unity (Art. 15). When a is less than 4^{-4} , or a less than about 0.33, the above series applies, and if ϵ designates one of the imaginary fifth roots of unity (Art. 15), the five roots of the equation are

$$\begin{aligned} y_1 = & 1 + a - a^2 + a^3 - \frac{2}{5}a^5 + \frac{7}{5}a^7 - \frac{1}{5}a^9 + \frac{2}{5}a^{11} - \dots, \\ y_2 = & \epsilon + \epsilon^2 a - \epsilon^3 a^2 + \epsilon^4 a^3 - \frac{2}{5}\epsilon a^5 + \frac{7}{5}\epsilon^2 a^7 - \frac{1}{5}\epsilon^3 a^9 + \frac{2}{5}\epsilon^4 a^{11} - \dots, \\ y_3 = & \epsilon^2 + \epsilon^4 a - \epsilon a^2 + \epsilon^3 a^3 - \frac{2}{5}\epsilon^2 a^5 + \frac{7}{5}\epsilon^4 a^7 - \frac{1}{5}\epsilon^3 a^9 + \frac{2}{5}\epsilon^4 a^{11} - \dots, \\ y_4 = & \epsilon^3 + \epsilon a - \epsilon^4 a^2 + \epsilon^2 a^3 - \frac{2}{5}\epsilon^3 a^5 + \frac{7}{5}\epsilon a^7 - \frac{1}{5}\epsilon^4 a^9 + \frac{2}{5}\epsilon^2 a^{11} - \dots, \\ y_5 = & \epsilon^4 + \epsilon^3 a - \epsilon^2 a^2 + \epsilon a^3 - \frac{2}{5}\epsilon^4 a^5 + \frac{7}{5}\epsilon^3 a^7 - \frac{1}{5}\epsilon^2 a^9 + \frac{2}{5}\epsilon^3 a^{11} - \dots \end{aligned}$$

For example, let $a=0.1$, or $y^5 - \frac{1}{2}y - 1 = 0$; then the value of y_1 is found to be $+1.09097$, while the other roots are

$$\begin{aligned} y_2 &= +0.23649 + 1.01470i, & y_3 &= -0.781975 + 0.48372i, \\ y_4 &= +0.23649 - 1.01470i, & y_5 &= -0.781975 - 0.48372i, \end{aligned}$$

which are correct in the fifth decimal place.

For the case where a^n is greater than $k^{-k}(n-k)^{k-n}b^n$ in the trinomial equation $y^n - n a y^{n-k} - b = 0$, the roots may be obtained by inserting x in other terms than the second. To illustrate the method by the quintic $y^5 - 5ay - 1 = 0$, let x be placed in the last term, giving $y^5 - 5ay - x = 0$; obtaining the derivatives and making $n=0$, there is found a series giving four of the roots, since $(5a)^{\frac{1}{4}}$ in this series has four values. Again, placing x in the first term the equation is $xy^5 - 5ay - 1 = 0$; and applying the method, there is found a series which gives the other root. It may also be shown that these series are convergent when a^5 is numerically greater than 4^{-4} . When $a^5 = 4^{-4}$ the quintic has two equal roots and the series do not apply, but in this case the equal roots are readily found (Art. 5) and after their removal the other three roots are found by the solution of a cubic equation.

When this method is applied to an algebraic equation of the n^{th} degree which contains more terms than three, there may be obtained several series by inserting x in different terms, and the series desired are those which are convergent. A general rule for selecting the terms which are to contain x is given by Lambert, and he applies the method to the solution of the quintic equation $y^5 - 10y^3 + 6y + 1 = 0$. First, writing $y^5 - 10y^3 + 6xy + x = 0$, the values of y_0 are $+3.167$ and -3.167 , those of $(dy/dx)_0$ are -1.00 and $+0.090$, and those of (d^2y/dx^2) are -0.016 and $+0.016$; inserting these in Maclaurin's formula there are found $y_1 = +3.05$ and $y_2 = -3.06$. Secondly, writing $xy^5 - 10y^3 + 6y + x = 0$, a series results which gives $y_3 = +0.87$ and $y_4 = -0.69$. Lastly, writing $xy^5 - 10xy^3 + 6y + 1$, there is found $y_5 = -0.17$.

This method may likewise be used for computing one of the roots of a transcendental equation, provided the resulting

series is convergent. For example, take $2y + \log y - 10000 = 0$. Writing $2y + x \log y - 10000 = 0$, there are found the values $y_0 = +5000$, $(dy/dx)_0 = -\frac{1}{2} \log y_0$, and $(d^2y/dx^2)_0 = +0.0001 \log y_0$. When the logarithm is in the common system the root is $y = 4998.15$; when it is in the Naperian system the root is $y = 4995.74$.

Prob. 24. Compute the roots of $x^3 - 2x - 2 = 0$ by the above method and also by that of Art. 9.

Prob. 25. The equation $y^4 - 11727y + 40385 = 0$ occurs in a paper on the precession of a viscous spheroid by G. H. Darwin in *Philosophical Transactions of the Royal Society*, 1879, Part ii, p. 508. Compute the four roots to five significant figures.

ART. 17. SYMMETRIC FUNCTIONS OF ROOTS.

The coefficients of an algebraic equation are the simplest symmetric functions of its roots. Let the equation be

$$x^n - ax^{n-1} + bx^{n-2} - cx^{n-3} + dx^{n-4} - \dots = 0, \quad \dots \quad (1)$$

and let x_1, x_2, x_3, \dots be its n roots. Then

$$\begin{aligned} a &= x_1 + x_2 + x_3 + \dots, & b &= x_1x_2 + x_2x_3 + x_3x_4 + \dots, \\ c &= x_1x_2x_3 + x_2x_3x_4 + \dots, & d &= x_1x_2x_3x_4 + x_2x_3x_4x_5 + \dots, \end{aligned}$$

and the last term is $\pm x_1x_2x_3 \dots x_n$. All symmetric functions of the roots may be expressed in terms of the coefficients.

The sums of the powers of the roots are important symmetric functions. Let S_m represent $x_1^m + x_2^m + x_3^m + \dots$; then when m is equal to or less than n , the following are the Newtonian expressions for the sums of the powers of the roots:

$$\begin{aligned} S_1 &= a, & S_2 &= a^2 - 2b, & S_3 &= a^3 - 3ab + 3c, \\ S_4 &= a^4 - 4a^2b + 4ac + 2b^2 - 4d, & & \dots \end{aligned}$$

Let $\pm l$ represent the coefficient of the $(m+1)^{\text{th}}$ term in the general equation (1), this being $+$ when m is even and $-$ when m is odd. Then the following general formulas furnish values of S_m for all cases:

$$\begin{aligned} S_m - aS_{m-1} + bS_{m-2} - cS_{m-3} + \dots \pm ml &= 0, & m &\leq n, \\ S_{n+m} - aS_{n+m-1} + bS_{n+m-2} - \dots \pm lS_m &= 0, & m &> n. \end{aligned}$$

For example, take $x^3 - 2x - 2 = 0$, for which $a = 0$, $b = -2$, $c = +2$; then from the first formula $S_1 = 0$, $S_2 = 4$, $S_3 = 6$, and from the second formula $S_4 = 8$, $S_5 = 20$, $S_6 = 28$, etc.

Other important symmetric functions of the roots are the sums of the squares of the terms in the above expressions for the coefficients b , c , d , etc. Let these be called B , C , D , etc., or

$$B = x_1^2 x_2^2 + x_2^2 x_3^2 + \dots, \quad C = x_1^2 x_2^2 x_3^2 + x_2^2 x_3^2 x_4^2 + \dots,$$

and let it be required to find the values of B , C , D , etc., in terms of a , b , c , etc. For this purpose let (1) be written

$$x^n + bx^{n-2} + dx^{n-4} + \dots = ax^{n-1} + cx^{n-3} + ex^{n-5} + \dots,$$

and let both members be squared and the resulting equation be reduced to the form

$$y^n - Ay^{n-1} + By^{n-2} - Cy^{n-3} + Dy^{n-4} - \dots = 0, \quad \dots \quad (2)$$

in which y represents x^2 . This equation has n roots x_1^2 , x_2^2 , x_3^2 , \dots ; hence the value of A is $x_1^2 + x_2^2 + x_3^2 + \dots$, and the values of B and C are the symmetric functions above written. The algebraic work shows that

$$A = a^2 - 2b, \quad B = b^2 - 2ac + ad, \quad C = c^2 - 2bd + 2ae - 2f, \quad \dots,$$

and thus in general any coefficient in (2) is obtained from those in (1) by the following rule: the coefficient of y^m in (2) is found by taking the square of the coefficient of x^m in (1) together with twice the products of the coefficients of the terms equally removed from it to right and left, these products being alternately negative and positive.

An equation whose roots are the squares of those of (2) may be obtained by a similar process, the equation being

$$z^n - A_1 z^{n-1} + B_1 z^{n-2} - C_1 z^{n-3} + D_1 z^{n-4} - \dots = 0, \quad \dots \quad (3)$$

in which A_1 , B_1 , C_1 , \dots are computed from A , B , C , in the same manner that A , B , C , \dots were computed from (1). For example, take the equation $x^7 + 3x^4 + 6 = 0$; the equation whose roots are

squares of those of the given equation is $y^7 + 9y^4 + 36y^2 + 36 = 0$, and that whose roots are the fourth powers of those of the given equation is $z^7 + 81z^4 - 648z^3 + 1944z^2 - 2592z + 1296 = 0$.

Prob. 26. Find an equation the roots of which are the fourth powers of the roots of $x^3 + x + 10 = 0$.

Prob. 27. For the cubic equation $x^3 - ax^2 + bx - c = 0$ show that the value of $x_1^3 x_2^3 + x_2^3 x_3^3 + x_3^3 x_1^3$ is $b^3 - 3abc + 3c^2$.

Prob. 28. For the quartic equation $x^4 - ax^3 + bx^2 - cx + d = 0$ show that the value of S_5 is $a^5 - 5a^3b - 5ab^2 + 5a^2c - 5ad - 5bc$.

ART. 18. LOGARITHMIC SOLUTIONS.

A logarithmic method for the solution of algebraic equations with numerical coefficients was published by Gräffe in 1837 and exemplified by Encke in 1841.* The method involves the formation of an equation whose roots are high powers of the roots of the given equation; to do this an equation is first derived, by help of the principles in Art. 17, whose roots are the squares of those of the given equation, then one whose roots are the squares of those of the second equations or the fourth powers of those of the given equation, and so on. With the use of addition and subtraction logarithms, the greater part of the numerical work may be made logarithmic. The method is of especial value when all the roots of the given equation are real and unequal.

To illustrate the theory of the method, let p, q, r, s , etc., denote the roots, each of which is supposed to be a real negative number; let $[p]$ denote $p + q + r + \dots$, $[pq]$ denote $pq + qr + rs + \dots$, and so on. Then the general algebraic equation may be written

$$x^n - [p]x^{n-1} + [pq]x^{n-2} - [pqr]x^{n-3} + [pqrs]x^{n-4} - \dots, \quad (1)$$

and the equation whose roots are p^2, q^2, r^2, \dots is, by Art. 17,

$$y^n - [p^2]y^{n-1} + [p^2q^2]y^{n-2} - [p^2q^2r^2]y^{n-3} + [p^2q^2r^2s^2]y^{n-4} - \dots,$$

in which $[p^2]$ denotes $p^2 + q^2 + r^2 + \dots$, $[p^2q^2]$ denotes $p^2q^2 + q^2r^2 + \dots$,

* Crelle's Journal für Mathematik, 1841, Vol. XXII, pp. 193-248.

and so on. From this equation another may be derived having the roots p^4, q^4, r^4, \dots , and then another may be found having the roots p^8, q^8, r^8, \dots . This process can be continued until an equation is derived whose roots are p^m, q^m, r^m, \dots , where m is a power of 2 sufficiently high for the subsequent operations. This equation is

$$z^n - [p^m]z^{n-1} + [p^mq^m]z^{n-2} - [p^mq^mr^m]z^{n-3} + \dots$$

Now let p be the root of (1) which is largest in numerical value, q the next, r the next, and so on. Then, as m increases the value of $[p^m]$ approaches p^m , that of $[p^mq^m]$ approaches p^mq^m , that of $[p^mq^mr^m]$ approaches $p^mq^mr^m$, and so on. Hence when m is large $[p^m]$ is an approximation to the value of p^m , and $[p^mq^m]/[p^m]$ is an approximation to the value of q^m . Accordingly by making m sufficiently large, the values of p^m, q^m, r^m, \dots , and hence those of p, q, r, \dots , may be obtained to any required degree of numerical precision. When two roots are nearly equal numerically, it will be necessary to make m very large; when equal roots exist they should be removed by the usual method.

To illustrate the application of the method, let it be required to find the roots of the quintic equation

$$x^5 + 13x^4 - 81x^3 - 34x^2 + 464x - 181 = 0.$$

By comparison with (1) of Art. 17 it is seen that $a = -13, b = -81, c = +34, d = +464, e = +181$. The equation whose roots are the squares of those of the given quintic is now found from (2) of Art. 17, by computing $A = a^2 - 2b = 331, B = b^2 - 2ac + 2d = 8373, C = c^2 - 2bd + 2ae = 71618, D = d^2 - 2ce = 202988, E = e^2 = 32761$, and then

$$y^5 - 331y^4 + 8373y^3 - 71618y^2 + 202988y - 32761 = 0.$$

Taking the logarithms of the coefficients, this equation may be written

$$y^5 - (2.51983)y^4 + (3.92288)y^3 - (4.85502)y^2 + (5.30747)y - (4.51536) = 0,$$

in which the coefficients are expressed by their logarithms inclosed in parentheses. The logarithms of the coefficients for the equation whose roots are the fourth powers of the given quintic are now found by the use of addition and subtraction logarithmic tables, and this equation is

$$z^5 - (4.96762)z^4 + (7.36364)z^3 - (9.24342)z^2 + (10.56243)z - (9.03072) = 0.$$

Next the equation whose roots are the eighth powers of the roots of the given quintic is derived from the preceding one in a similar manner and is found to be

$$w^5 - (9.93290)w^4 + (14.31934)w^3 - (18.14025)w^2 + (21.12363)w - (18.06144) = 0,$$

and then the equation whose roots are the sixteenth powers of the roots of the given quintic is

$$v^5 - (19.86580)v^4 + (28.29778)v^3 - (36.13131)v^2 + (42.24726)v - (36.12288) = 0.$$

It is now observed that the coefficients of the second, fourth, and fifth terms in the equation for v are the squares of those of the similar terms in the equation for w . Hence two of the roots are now determined as follows:

$$\begin{aligned} \log p^8 &= 9.93290, & \log p &= 1.24161, & p &= 17.443; \\ \log t^8 &= 18.06144 - 21.12363, & \log t &= 1.61723, & t &= 0.4142. \end{aligned}$$

These are the numerical values of the largest and smallest roots of the given quintic, but the method does not determine whether they are positive or negative; by trial in the given quintic it will be found that -17.443 and $+0.4142$ are roots. To obtain the others, the process must be continued until two successive equations are found for which all the coefficients in the second are the squares of those in the first. Since in this case two roots lie near together, the process does not terminate, with five-place logarithms, until the 512^{th} powers are reached. The three

remaining roots are thus found to be $q = +3.230$, $r = +3.213$, and $s = -1.4142$.

When this method is applied to an algebraic equation which has imaginary roots, this fact is indicated by the deviation of signs of the terms in the power equations from the form as given in (2) of Art. 17; that is, these signs are not alternately positive and negative. As an example of such a case Encke applies the process to the equation

$$x^7 - 2x^5 - 3x^3 + 4x^2 - 5x + 6 = 0,$$

and deduces for the equation of the 256th powers of the roots

$$v^7 - (74.95884)v^6 + (122.81202)v^5 + (151.32153)v^4 + (179.58882)v^3 - (190.99129)v^2 - (195.21132)v - (199.20704) = 0.$$

Here it is seen that the coefficients of v^4 and v have signs opposite to those of the normal form, and hence two pairs of imaginary roots are indicated. The real roots of the given equation are then determined as follows:

$$\begin{aligned} \log x_1^{256} &= 74.95884, & \log x_1 &= 0.29281, & x_1 &= -1.9625, \\ \log x_2^{256} &= 122.81202 - 74.95886, & \log x_2 &= 0.18693, & x_2 &= +1.5379, \\ \log x_6^{256} &= 190.99129 - 179.58882, & \log x_6 &= 0.04454, & x_6 &= +1.1080, \end{aligned}$$

while the logarithms of the moduli of the imaginary pairs may be obtained by taking the difference of the logarithms of v^5 and v^3 and that of v^2 and v^0 , and dividing each by 512. It is then not difficult to show that the two quadratic equations

$$x^2 - 0.60921x + 1.07668 = 0, \quad x^2 + 1.29263 + 1.66642i = 0,$$

furnish the imaginary roots of the given equation of the seventh degree.

Prob. 29. Compute the roots of $x^5 - 10x^3 + 6x + 1 = 0$.

Prob. 30. How many real roots has the equation $x^7 + 3x^4 + 6 = 0$? Can they be advantageously computed by the above method? What is the best method for finding the roots to four decimal places?

ART. 19. INFINITE EQUATIONS.

An infinite series containing ascending powers of x may be equated to zero and be called an infinite equation. For example, consider the equation

$$x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \frac{x^9}{9!} - \dots = 0,$$

in which the first member is the expansion of $\sin x$; this equation has the roots $0, \pi, 2\pi, 3\pi$, etc., since these are the values which satisfy the equation $\sin x = 0$. Again,

$$1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \frac{x^8}{8!} + \dots = 0$$

is the same as $\cosh x = 0$, and hence its roots are $\frac{1}{2}\pi i, \frac{3}{2}\pi i$, etc.

The series known as Bessel's first function when equated to zero furnishes an infinite equation whose roots are of interest in the theory of heat*; this equation is

$$1 - \frac{x^2}{2^2} + \frac{x^4}{2^2 \cdot 4^2} - \frac{x^6}{2^2 \cdot 4^2 \cdot 6^2} + \frac{x^8}{2^2 \cdot 4^2 \cdot 6^2 \cdot 8^2} - \dots = 0,$$

and it has an infinite number of real positive roots, the smallest of which is 2.4048. The roots of equations of this kind may be computed by tentative methods, and when they are approximately known Newton's rule (Art. 4) may be used to obtain more precise values.

As an example take another equation which also occurs in the theory of heat, namely,

$$1 - x + \frac{x^2}{(2!)^2} - \frac{x^3}{(3!)^2} + \frac{x^4}{(4!)^2} - \frac{x^5}{(5!)^2} + \dots = 0.$$

It is plain that this equation can have no negative roots, for a negative value of x renders all the terms of the first member

* Mathematical Monograph, No. 5, pp. 23, 63.

positive. Calling the first member $f(x)$, the first derivative is

$$f'(x) = -1 + \frac{x}{2} - \frac{x^2}{2^2 \cdot 3} + \frac{x^3}{2^2 \cdot 3^2 \cdot 4} - \frac{x^4}{2^2 \cdot 3^2 \cdot 4^2 \cdot 5}.$$

By trial it may be found that one root of $f(x)=0$ lies between 1.44 and 1.45. For $x=1.44$, $f(x)$ becomes $+0.002508$ and $f'(x)$ becomes $+0.4334$. Then $f(x)/f'(x)=0.0058$, and accordingly $x_1=1.44+0.0058=1.4458$ is one of the roots. Another root of this equation is $x_2=7.6178$. In general equations of this kind have an infinite number of roots.

The term infinite is sometimes applied to an algebraic equation having an infinite root, and cases of this kind are often stated as curious mathematical problems. For instance, the solution of the equation

$$x-a=(x^2-a\sqrt{x^2+a^2})^{\frac{1}{2}},$$

when made by squaring each member twice, gives the roots $x=\frac{4}{3}a$ and $x=0$. But $x=0$ does not satisfy the equation as written, although it applies if the sign of the second radical be changed. The equation, however, may be put in the form

$$1-\frac{a}{x}=\left(1-\sqrt{\frac{a^2}{x^2}+\frac{a^4}{x^4}}\right)^{\frac{1}{2}},$$

and it is now seen that $x=\infty$ is one of its roots. The false value $x=0$ arises from the circumstance that the squaring operations give results which may be also derived from equations having signs before the radicals different from those written in the given equation.

Prob. 31. Differentiate the above function of Bessel and equate the derivative to zero. Compute two of the roots of this infinite equation.

Prob. 32. Find the roots of $2\sqrt{x-2}=\sqrt{x-3}+\sqrt{x-1}$.

Prob. 33. Consult a paper by Stern in Crelle's Journal für Mathematik, 1841, pp. 1-62, and explain his methods of solving the equations $\cos x \cosh x + 1 = 0$ and $(4-3x^2) \sin x - 4x \cos x = 0$.

ART. 20. NOTES AND PROBLEMS.

The algebraic solutions of the quadratic, cubic, and quartic equations are valid for imaginary coefficients also. In general the roots of such equations are all imaginary. The method of McClintock (Art. 14) and that of Lambert (Art. 16) may also be applied to the expression of the roots of these equations in infinite series.

As an illustration take the equation $x^3 - 3x + 4i = 0$. By any method may be found the roots $x_1 = -i$, $x_2 = -0.5i + 1.936$ and $x_3 = -0.5i - 1.936$; two of the roots here form a pair in which the imaginary part is the same for both, the real and imaginary parts of the complex quantities having changed places. There are, however, many equations with imaginary and complex coefficients in which pairs of roots do not occur.

The most general case of an algebraic equation is when the coefficients a, b, c, \dots in (1) of Art. 17 are complex quantities of the form $m + ni$, $p + qi$, \dots . Such equations rarely, if ever, occur in physical investigations, but the general methods explained in the preceding pages will usually suffice for their solution, approximate values of the roots being first obtained by trial if necessary. In general the roots of such equations are all complex, although conditions between m and n , p and q , etc., may be introduced which will render real one or more of the roots.

Prob. 34. Show that the equation $x - e^x = 0$ has many pairs of imaginary roots and that the smallest roots are $0.3181 \pm 1.3372i$.

Prob. 35. Solve $\frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \dots = -1$.

Prob. 36. Discuss the equation $x - \tan x = 0$ and show that its smallest root is 4.49341 .

Prob. 37. Find the value of x in the equation $e^{\pi x} + 1 = 0$, and also that in the equation $e^{\frac{1}{2}\pi x} - i = 0$.

Prob. 38. Show that $x^2 + (a + bi)x + c + di = 0$ has one real and one complex root when the coefficients are so related that $b^2c + d^2 - abd = 0$.

Prob. 39. When and by whom was the sign of equality first used? What reason was given as to the propriety of its use for this purpose?

Prob. 40. There is a conical glass, 6 inches deep, and the diameter at the top is 5 inches. When it is one-fifth full of water, a sphere 4 inches in diameter is put into the glass. What part of the vertical diameter of the sphere is immersed in the water?

Prob. 41. When seven ordinates are to be erected upon an abscissa line of unit length in order to determine the area between that line and a curve, their distances apart in order to give the most advantageous result are, according to Gauss, determined by the equation

$$x^7 - \frac{7}{2}x^6 + \frac{63}{13}x^5 - \frac{175}{52}x^4 + \frac{175}{143}x^3 - \frac{63}{286}x^2 + \frac{7}{429}x - \frac{1}{3432} = 0.$$

Compute the roots to five decimal places and compare them with those given by Gauss.

INDEX.

- Abel's discussion of quintic, 22.
Algebraic equations, 1, 2.
 solutions, 15-24.
Approximation of roots, 3, 12, 49.
 rule, 6.
Bessel's function, 43.
Binomial equations, 16, 26, 31.
Cardan's formula, 17, 18, 28, 29.
Catenary, 14.
Cube roots of unity, 32.
Cubic equations, 3, 17, 28.
Cylinder, floating, 13.
De Moivre's quintic, 22, 26.
 theorem, 33.
Derivative equation, 9.
Elliptic solution of quintic, 23.
Fifth roots of unity, 16, 32.
Graphic solutions, 3.
Graphs of equations, 9.
Gräffe's method, 14.
Horner's process, 2, 12.
Howe truss strut problem, 21.
Hudde's method, 8, 12.
Imaginary coefficients, 45.
 roots, 11, 18, 20, 30, 34, 42.
Infinite equations, 43.
Lagrange's resolvent, 15.
Lambert's method, 33.
Literal equations, 1, 10.
Logarithmic solutions, 39.
Maclaurin's formula, 34.
McClintock's quintic discussion, 23.
 series method, 29.
Newton's approximation rule, 6.
Numerical equations, 1, 10.
Powers of roots, 38, 40.
Properties of equations, 11.
Quadratic equations, 16.
Quartic equations, 19, 20.
Quintic equations, 21, 36.
Real roots, 2, 3, 12, 40.
Regula falsi, 5.
Removal of terms, 22.
Resolvent, 17.
Root, 1.
Roots in series, 27, 29, 31, 34.
 of unity, 16, 31.
Separation of roots, 8.
Sixth roots of unity, 17, 32.
Sphere, floating, 13, 25.
Sturm's theorem, 8, 12.
Symmetric functions, 37.
Transcendental equations, 2, 4, 13.
Trigonometric solutions, 24, 26.
Trinomial equations, 29, 35.
Tschirnhausen's transformation, 22.
Vectors, 16, 33.
Water-pipe problem, 13.

SHORT-TITLE CATALOGUE

OF THE

PUBLICATIONS

OF

JOHN WILEY & SONS

NEW YORK

LONDON: CHAPMAN & HALL, LIMITED

ARRANGED UNDER SUBJECTS

Descriptive circulars sent on application. Books marked with an asterisk (*) are sold at *net* prices only. All books are bound in cloth unless otherwise stated.

AGRICULTURE—HORTICULTURE—FORESTRY.

Armsby's Principles of Animal Nutrition.....	8vo,	\$4 00
Budd and Hansen's American Horticultural Manual:		
Part I. Propagation, Culture, and Improvement.....	12mo,	1 50
Part II. Systematic Pomology.....	12mo,	1 50
Elliott's Engineering for Land Drainage.....	12mo,	1 50
Practical Farm Drainage. (Second Edition, Rewritten.).....	12mo,	1 50
Graves's Forest Mensuration.....	8vo,	4 00
Green's Principles of American Forestry.....	12mo,	1 50
Grotenfelt's Principles of Modern Dairy Practice. (Woll.).....	12mo,	2 00
* Herrick's Denatured or Industrial Alcohol.....	8vo,	4 00
Kemp and Waugh's Landscape Gardening. (New Edition, Rewritten. In Preparation.)		
* McKay and Larsen's Principles and Practice of Butter-making.....	8vo,	1 50
Maynard's Landscape Gardening as Applied to Home Decoration.....	12mo,	1 50
Sanderson's Insects Injurious to Staple Crops.....	12mo,	1 50
Sanderson and Headlee's Insects Injurious to Garden Crops. (In Preparation.)		
* Schwarz's Longleaf Pine in Virgin Forest.....	12mo,	1 25
Stockbridge's Rocks and Soils.....	8vo,	2 50
Winton's Microscopy of Vegetable Foods.....	8vo,	7 50
Woll's Handbook for Farmers and Dairymen.....	16mo,	1 50

ARCHITECTURE.

Baldwin's Steam Heating for Buildings.....	12mo,	2 50
Berg's Buildings and Structures of American Railroads.....	4to,	5 00
Birkmire's Architectural Iron and Steel.....	8vo,	3 50
Compound Riveted Girders as Applied in Buildings.....	8vo,	2 00
Planning and Construction of American Theatres.....	8vo,	3 00
Planning and Construction of High Office Buildings.....	8vo,	3 50
Skeleton Construction in Buildings.....	8vo,	3 00

Briggs's Modern American School Buildings.....	8vo,	\$4 00
Byrne's Inspection of Materials and Workmanship Employed in Construction.....	16mo,	3 00
Carpenter's Heating and Ventilating of Buildings.....	8vo,	4 00
* Corthell's Allowable Pressure on Deep Foundations.....	12mo,	1 25
Freitag's Architectural Engineering.....	8vo,	3 50
Fireproofing of Steel Buildings.....	8vo,	2 50
Gerhard's Guide to Sanitary Inspections. (Fourth Edition, Entirely Revised and Enlarged.).....	12mo,	1 50
* Modern Baths and Bath Houses.....	8vo,	3 00
Sanitation of Public Buildings.....	12mo,	1 50
Theatre Fires and Panics.....	12mo,	1 50
* The Water Supply, Sewerage and Plumbing of Modern City Buildings.....	8vo,	4 00
Johnson's Statics by Algebraic and Graphic Methods.....	8vo,	2 00
Kellaway's How to Lay Out Suburban Home Grounds.....	8vo,	2 00
Kidder's Architects' and Builders' Pocket-book.....	16mo, mor.,	5 00
Merrill's Stones for Building and Decoration.....	8vo,	5 00
Monckton's Stair-building.....	4to,	4 00
Patton's Practical Treatise on Foundations.....	8vo,	5 00
Peabody's Naval Architecture.....	8vc,	7 50
Rice's Concrete-block Manufacture.....	8vo,	2 00
Richey's Handbook for Superintendents of Construction.....	16mo, mor.	4 00
Building Foreman's Pocket Book and Ready Reference..	16mo, mor.	5 00
* Building Mechanics' Ready Reference Series:		
* Carpenters' and Woodworkers' Edition.....	16mo, mor.	1 50
* Cement Workers' and Plasterers' Edition.....	16mo, mor.	1 50
* Plumbers', Steam-Fitters', and Tinnners' Edition..	16mo, mor.	1 50
* Stone- and Brick-masons' Edition.....	16mo, mor.	1 50
Sabin's House Painting.....	12mo,	1 00
Siebert and Biggin's Modern Stone-cutting and Masonry.....	8vo,	1 50
Snow's Principal Species of Wood.....	8vo,	3 50
Towne's Locks and Builders' Hardware.....	16mo, mor.	3 00
Wait's Engineering and Architectural Jurisprudence.....	8vo,	6 00
Law of Contracts.....	8vo,	3 00
Law of Operations Preliminary to Construction in Engineering and Architecture.....	8vo,	5 00
Sheep,		5 50
Wilson's Air Conditioning.....	12mo,	1 50
Worcester and Atkinson's Small Hospitals, Establishment and Maintenance, Suggestions for Hospital Architecture, with Plans for a Small Hospital.....	12mo,	1 25

ARMY AND NAVY.

Bernadou's Smokeless Powder, Nitro-cellulose, and the Theory of the Cellulose Molecule.....	12mo,	2 50
Chase's Art of Pattern Making.....	12mo,	2 50
Screw Propellers and Marine Propulsion.....	8vo,	3 00
* Cloke's Enlisted Specialists' Examiner.....	8vo,	2 00
* Gunner's Examiner.....	8vo,	1 50
Craig's Azimuth.....	4to,	3 50
Crehore and Squier's Polarizing Photo-chronograph.....	8vo,	3 50
* Davis's Elements of Law.....	8vo,	2 50
* Treatise on the Military Law of United States.....	8vo,	7 00
* Dudley's Military Law and the Procedure of Courts-martial..	Large 12mo,	2 50
Durand's Resistance and Propulsion of Ships.....	8vo,	5 00
* Dyer's Handbook of Light Artillery.....	12mo,	3 00
Eissler's Modern High Explosives.....	8vo,	4 00
* Fieberger's Text-book on Field Fortification.....	Large 12mo,	2 00
Hamilton and Bond's The Gunner's Catechism.....	18mo,	1 00
* Hoff's Elementary Naval Tactics.....	8vo,	1 50

Ingalls's Handbook of Problems in Direct Fire.....	8vo,	\$4 00
* Lissak's Ordnance and Gunnery.....	8vo,	6 00
* Ludlow's Logarithmic and Trigonometric Tables.....	8vo,	1 00
* Lyons's Treatise on Electromagnetic Phenomena. Vols. I. and II.....	8vo, each,	6 00
* Mahan's Permanent Fortifications. (Mercur.).....	8vo, half mor.	7 50
Manual for Courts-martial.....	16mo, mor.	1 50
* Mercur's Attack of Fortified Places.....	12mo,	2 00
* Elements of the Art of War.....	8vo,	4 00
Nixon's Adjutants' Manual.....	24mo,	1 00
Peabody's Naval Architecture.....	8vo,	7 50
* Phelps's Practical Marine Surveying.....	8vo,	2 50
Putnam's Nautical Charts.....	8vo,	2 00
Rust's Ex-meridian Altitude, Azimuth and Star-Finding Tables.....	8vo,	5 00
Selkirk's Catechism of Manual of Guard Duty. (In Press.)		
Sharpe's Art of Subsisting Armies in War.....	18mo, mor.	1 50
Taylor's Speed and Power of Ships. (In Press.)		
* Tupes and Poole's Manual of Bayonet Exercises and Musketry Fencing.		
	24mo, leather,	50
* Weaver's Military Explosives.....	8vo,	3 00
* Woodhull's Military Hygiene for Officers of the Line.....	Large 12mo,	1 50

ASSAYING.

Betts's Lead Refining by Electrolysis.....	8vo,	4 00
Fletcher's Practical Instructions in Quantitative Assaying with the Blowpipe.		
	16mo, mor.	1 50
Furman and Pardoe's Manual of Practical Assaying. (Sixth Edition, Revised and Enlarged.).....	8vo,	3 00
Lodge's Notes on Assaying and Metallurgical Laboratory Experiments.....	8vo,	3 00
Low's Technical Methods of Ore Analysis.....	8vo,	3 00
Miller's Cyanide Process.....	12mo,	1 00
Manual of Assaying.....	12mo,	1 00
Minet's Production of Aluminum and its Industrial Use. (Waldo.).....	12mo,	2 50
Ricketts and Miller's Notes on Assaying.....	8vo,	3 00
Robine and Lenglen's Cyanide Industry. (Le Clerc.).....	8vo,	4 00
Seamon's Manual for Assayers and Chemists. (In Press.)		
Ulke's Modern Electrolytic Copper Refining.....	8vo,	3 00
Wilson's Chlorination Process.....	12mo,	1 50
Cyanide Processes.....	12mo,	1 50

ASTRONOMY.

Comstock's Field Astronomy for Engineers.....	8vo,	2 50
Craig's Azimuth.....	4to,	3 50
Crandall's Text-book on Geodesy and Least Squares.....	8vo,	3 00
Doolittle's Treatise on Practical Astronomy.....	8vo,	4 00
Hayford's Text-book of Geodetic Astronomy.....	8vo,	3 00
Hosmer's Azimuth.....	16mo, mor.	1 00
Merriman's Elements of Precise Surveying and Geodesy.....	8vo,	2 50
* Michie and Harlow's Practical Astronomy.....	8vo,	3 00
Rust's Ex-meridian Altitude, Azimuth and Star-Finding Tables.....	8vo,	5 00
* White's Elements of Theoretical and Descriptive Astronomy.....	12mo,	2 00

CHEMISTRY.

* Abderhalden's Physiological Chemistry in Thirty Lectures. (Hall and Defren.).....	8vo,	5 00
* Abegg's Theory of Electrolytic Dissociation. (von Ende.).....	12mo,	1 25
Alexeyeff's General Principles of Organic Syntheses. (Matthews.).....	8vo,	3 00
Allen's Tables for Iron Analysis.....	8vo,	3 00
Armsby's Principles of Animal Nutrition.....	8vo,	4 00
Arnold's Compendium of Chemistry. (Mandel.).....	Large 12mo,	3 50

Association of State and National Food and Dairy Departments, Hartford Meeting, 1906.....	8vo,	\$3 00
Jamestown Meeting, 1907.....	8vo,	3 00
Austen's Notes for Chemical Students.....	12mo,	1 50
Baskerville's Chemical Elements. (In Preparation.)		
Bernadou's Smokeless Powder.—Nitro-cellulose, and Theory of the Cellulose Molecule.....	12mo,	2 50
* Biltz's Introduction to Inorganic Chemistry. (Hall and Phelan.)	12mo,	1 25
Laboratory Methods of Inorganic Chemistry. (Hall and Blanchard.)	8vo,	3 00
* Blanchard's Synthetic Inorganic Chemistry.....	12mo,	1 00
* Browning's Introduction to the Rarer Elements.....	8vo,	1 50
* Claassen's Beet-sugar Manufacture. (Hall and Rolfe.)	8vo,	3 00
Classen's Quantitative Chemical Analysis by Electrolysis. (Boltwood.)	8vo,	3 00
Cohn's Indicators and Test-papers. (Prescott.)	12mo,	2 00
Tests and Reagents.....	8vo,	3 00
* Danneel's Electrochemistry. (Merriam.)	12mo,	1 25
Dannerth's Methods of Textile Chemistry.....	12mo,	2 00
Duhem's Thermodynamics and Chemistry. (Burgess.)	8vo,	4 00
Effront's Enzymes and their Applications. (Prescott.)	8vo,	3 00
Eissler's Modern High Explosives.....	8vo,	4 00
Erdmann's Introduction to Chemical Preparations. (Dunlap.)	12mo,	1 25
* Fischer's Physiology of Alimentation.....	Large 12mo,	2 00
Fletcher's Practical Instructions in Quantitative Assaying with the Blowpipe.	16mo, mor.	1 50
Fowler's Sewage Works Analyses.....	12mo,	2 00
Fresenius's Manual of Qualitative Chemical Analysis. (Wells.)	8vo,	5 00
Manual of Qualitative Chemical Analysis. Part I. Descriptive. (Wells.)	8vo,	3 00
Quantitative Chemical Analysis. (Cohn.) 2 vols.	8vo,	12 50
When Sold Separately, Vol. I, \$6. Vol. II, \$8.		
Fuertes's Water and Public Health.....	12mo,	1 50
Furman and Pardoe's Manual of Practical Assaying. (Sixth Edition, Revised and Enlarged.)	8vo,	3 00
* Getman's Exercises in Physical Chemistry.....	12mo,	2 00
Gill's Gas and Fuel Analysis for Engineers.....	12mo,	1 25
* Gooch and Browning's Outlines of Qualitative Chemical Analysis.	Large 12mo,	1 25
Grotenfelt's Principles of Modern Dairy Practice. (Woll.)	12mo,	2 00
Groth's Introduction to Chemical Crystallography (Marshall).	12mo,	1 25
Hammarsten's Text-book of Physiological Chemistry. (Mandel.)	8vo,	4 00
Hanausek's Microscopy of Technical Products. (Winton.)	8vo,	5 00
* Haskins and Macleod's Organic Chemistry.....	12mo,	2 00
Hering's Ready Reference Tables (Conversion Factors).....	16mo, mor.	2 50
* Herrick's Denatured or Industrial Alcohol.....	8vo,	4 00
Hinds's Inorganic Chemistry.....	8vo,	3 00
* Laboratory Manual for Students.....	12mo,	1 00
* Holleman's Laboratory Manual of Organic Chemistry for Beginners. (Walker.)	12mo,	1 00
Text-book of Inorganic Chemistry. (Cooper.)	8vo,	2 50
Text-book of Organic Chemistry. (Walker and Mott.)	8vo,	2 50
* Holley's Lead and Zinc Pigments.....	Large 12mo,	3 00
Holley and Ladd's Analysis of Mixed Paints, Color Pigments, and Varnishes.	Large 12mo,	2 50
Hopkins's Oil-chemists' Handbook.....	8vo,	3 00
Jackson's Directions for Laboratory Work in Physiological Chemistry.	8vo,	1 25
Johnson's Rapid Methods for the Chemical Analysis of Special Steels, Steel-making Alloys and Graphite.....	Large 12mo,	3 00
Landauer's Spectrum Analysis. (Tingle.)	8vo,	3 00
Lassar-Cohn's Application of Some General Reactions to Investigations in Organic Chemistry. (Tingle.)	12mo,	1 00
Leach's Inspection and Analysis of Food with Special Reference to State Control.....	8vo,	7 50
Löb's Electrochemistry of Organic Compounds. (Lorenz.)	8vo,	3 00
Lodge's Notes on Assaying and Metallurgical Laboratory Experiments.	8vo,	3 00
Low's Technical Method of Ore Analysis.....	8vo,	3 00
Lowe's Paint for Steel Structures.....	12mo,	1 00
Lunge's Techno-chemical Analysis. (Cohn.)	12mo,	1 00

* McKay and Larsen's Principles and Practice of Butter-making.....	8vo,	\$1 50
Maire's Modern Pigments and their Vehicles.....	12mo,	2 00
Mandel's Handbook for Bio-chemical Laboratory.....	12mo,	1 50
* Martin's Laboratory Guide to Qualitative Analysis with the Blowpipe	12mo,	0 60
Mason's Examination of Water. (Chemical and Bacteriological.).....	12mo,	1 25
Water-supply. (Considered Principally from a Sanitary Standpoint.)	8vo,	4 00
* Mathewson's First Principles of Chemical Theory.....	8vo,	1 00
Matthews's Laboratory Manual of Dyeing and Textile Chemistry.....	8vo,	3 50
Textile Fibres. 2d Edition, Rewritten.....	8vo,	4 00
* Meyer's Determination of Radicles in Carbon Compounds. (Tingle.)		
Third Edition.....	12mo,	1 25
Miller's Cyanide Process.....	12mo,	1 00
Manual of Assaying.....	12mo,	1 00
Minet's Production of Aluminum and its Industrial Use. (Waldo.).....	12mo,	2 50
* Mittelstaedt's Technical Calculations for Sugar Works. (Bourbakis.)	12mo,	1 50
Mixter's Elementary Text-book of Chemistry.....	12mo,	1 50
Morgan's Elements of Physical Chemistry.....	12mo,	3 00
Outline of the Theory of Solutions and its Results.....	12mo,	1 00
* Physical Chemistry for Electrical Engineers.....	12mo,	1 50
* Moore's Outlines of Organic Chemistry.....	12mo,	1 50
Morse's Calculations used in Cane-sugar Factories.....	16mo, mor.	1 50
* Muir's History of Chemical Theories and Laws.....	8vo,	4 00
Mulliken's General Method for the Identification of Pure Organic Compounds.		
Vol. I. Compounds of Carbon with Hydrogen and Oxygen. Large	8vo,	5 00
Vol. II. Nitrogenous Compounds. (In Preparation.)		
Vol. III. The Commercial Dyestuffs.....	Large 8vo,	5 00
* Nelson's Analysis of Drugs and Medicines.....	12mo,	3 00
O'Driscoll's Notes on the Treatment of Gold Ores.....	8vo,	2 00
Ostwald's Conversations on Chemistry. Part One. (Ramsey.).....	12mo,	1 50
Part Two. (Turnbull.).....	12mo,	2 00
Introduction to Chemistry. (Hall and Williams.) (In Preparation.)		
Owen and Standage's Dyeing and Cleaning of Textile Fabrics.....	12mo,	2 00
* Palmer's Practical Test Book of Chemistry.....	12mo,	1 00
* Pauli's Physical Chemistry in the Service of Medicine. (Fischer.).....	12mo,	1 25
Penfield's Tables of Minerals, Including the Use of Minerals and Statistics		
of Domestic Production.....	8vo,	1 00
Pictet's Alkaloids and their Chemical Constitution. (Biddle.).....	8vo,	5 00
Poole's Calorific Power of Fuels.....	8vo,	3 00
Prescott and Winslow's Elements of Water Bacteriology, with Special Refer-		
ence to Sanitary Water Analysis.....	12mo,	1 50
* Reisig's Guide to Piece-Dyeing.....	8vo,	25 00
Richards and Woodman's Air, Water, and Food from a Sanitary Stand-		
point.....	8vo,	2 00
Ricketts and Miller's Notes on Assaying.....	8vo,	3 00
Rideal's Disinfection and the Preservation of Food.....	8vo,	4 00
Sewage and the Bacterial Purification of Sewage.....	8vo,	4 00
Riggs's Elementary Manual for the Chemical Laboratory.....	8vo,	1 25
Robine and Lenglen's Cyanide Industry. (Le Clerc.).....	8vo,	4 00
Ruddiman's Incompatibilities in Prescriptions.....	8vo,	2 00
Whys in Pharmacy.....	12mo,	1 00
* Ruer's Elements of Metallography. (Mathewson.).....	8vo,	3 00
Sabin's Industrial and Artistic Technology of Paint and Varnish.....	8vo,	3 00
Salkowski's Physiological and Pathological Chemistry. (Orndorff.).....	8vo,	2 50
Schimpf's Essentials of Volumetric Analysis.....	12mo,	1 25
Manual of Volumetric Analysis. (Fifth Edition, Rewritten).....	8vo,	5 00
* Qualitative Chemical Analysis.....	8vo,	1 25
Seamon's Manual for Assayers and Chemists. (In Press.)		
Smith's Lecture Notes on Chemistry for Dental Students.....	8vo,	2 50
Spencer's Handbook for Cane Sugar Manufacturers.....	16mo, mor.	3 00
Handbook for Chemists of Beet-sugar Houses.....	16mo, mor.	3 00
Stockbridge's Rocks and Soils.....	8vo,	2 50
Stone's Practical Testing of Gas and Gas Meters.....	8vo,	3 50
* Tillman's Descriptive General Chemistry.....	8vo,	3 00
Elementary Lessons in Heat.....	8vo,	1 50
Treadwell's Qualitative Analysis. (Hall.).....	8vo,	3 00

Soper's Air and Ventilation of Subways.....	12mo,	\$2 50
* Tracy's Exercises in Surveying.....	12mo, mor.	1 00
Tracy's Plane Surveying.....	16mo, mor.	3 00
* Trautwine's Civil Engineer's Pocket-book.....	16mo, mor.	5 00
Venable's Garbage Crematories in America.....	8vo,	2 00
Methods and Devices for Bacterial Treatment of Sewage.....	8vo,	3 00
Wait's Engineering and Architectural Jurisprudence.....	8vo,	6 00
	Sheep,	6 50
Law of Contracts.....	8vo,	3 00
Law of Operations Preliminary to Construction in Engineering and Architecture.....	8vo,	5 00
	Sheep,	5 50
Warren's Stereotomy—Problems in Stone-cutting.....	8vo,	2 50
* Waterbury's Vest-Pocket Hand-book of Mathematics for Engineers. 2 $\frac{1}{4}$ × 5 $\frac{3}{8}$ inches, mor.		1 00
* Enlarged Edition, Including Tables.....	mor.	1 50
Webb's Problems in the Use and Adjustment of Engineering Instruments. 16mo, mor.		1 25
Wilson's Topographic Surveying.....	8vo,	3 50

BRIDGES AND ROOFS.

Boller's Practical Treatise on the Construction of Iron Highway Bridges.....	8vo,	2 00
* Thames River Bridge.....	Oblong paper,	5 00
Burr and Falk's Design and Construction of Metallic Bridges.....	8vo,	5 00
Influence Lines for Bridge and Roof Computations.....	8vo,	3 00
Du Bois's Mechanics of Engineering. Vol. II.....	Small 4to,	10 00
Foster's Treatise on Wooden Trestle Bridges.....	4to,	5 00
Fowler's Ordinary Foundations.....	8vo,	3 50
Greene's Arches in Wood, Iron, and Stone.....	8vo,	2 50
Bridge Trusses.....	8vo,	2 50
Roof Trusses.....	8vo,	1 25
Grimm's Secondary Stresses in Bridge Trusses.....	8vo,	2 50
Heller's Stresses in Structures and the Accompanying Deformations.....	8vo,	3 00
Howe's Design of Simple Roof-trusses in Wood and Steel.....	8vo,	2 00
Symmetrical Masonry Arches.....	8vo,	2 50
Treatise on Arches.....	8vo,	4 00
* Jacoby's Structural Details, or Elements of Design in Heavy Framing.....	8vo,	2 25
Johnson, Bryan and Turneure's Theory and Practice in the Designing of Modern Framed Structures.....	Small 4to,	10 00
* Johnson, Bryan and Turneure's Theory and Practice in the Designing of Modern Framed Structures. New Edition. Part I.....	8vo,	3 00
Merriman and Jacoby's Text-book on Roofs and Bridges:		
Part I. Stresses in Simple Trusses.....	8vo,	2 50
Part II. Graphic Statics.....	8vo,	2 50
Part III. Bridge Design.....	8vo,	2 50
Part IV. Higher Structures.....	8vo,	2 50
Morison's Memphis Bridge.....	Oblong 4to,	10 00
Sondericker's Graphic Statics, with Applications to Trusses, Beams, and Arches.....	8vo,	2 00
Waddell's De Pontibus, Pocket-book for Bridge Engineers.....	16mo, mor.	2 00
* Specifications for Steel Bridges.....	12mo,	50
Waddell and Harrington's Bridge Engineering. (In Preparation.)		
Wright's Designing of Draw-spans. Two parts in one volume.....	8vo,	3 50

HYDRAULICS.

Barnes's Ice Formation.....	8vo,	3 00
Bazin's Experiments upon the Contraction of the Liquid Vein Issuing from an Orifice. (Trautwine.).....	8vo,	2 00
Bovey's Treatise on Hydraulics.....	8vo,	5 00
Church's Diagrams of Mean Velocity of Water in Open Channels. Oblong 4to, paper,		1 50
Hydraulic Motors.....	8vo,	2 00

Coffin's Graphical Solution of Hydraulic Problems.....	16mo, mor.	\$2 50
Plather's Dynamometers, and the Measurement of Power.....	12mo,	3 00
Folwell's Water-supply Engineering.....	8vo,	4 00
Frizell's Water-power.....	8vo,	5 00
Fuertes's Water and Public Health.....	12mo,	1 50
Water-filtration Works.....	12mo,	2 50
Ganguillet and Kutter's General Formula for the Uniform Flow of Water in Rivers and Other Channels. (Hering and Trautwine.).....	8vo,	4 00
Hazen's Clean Water and How to Get It.....	Large 12mo,	1 50
Filtration of Public Water-supplies.....	8vo,	3 00
Hazelhurst's Towers and Tanks for Water-works.....	8vo,	2 50
Herschel's 115 Experiments on the Carrying Capacity of Large, Riveted, Metal Conduits.....	8vo,	2 00
Hoyt and Grover's River Discharge.....	8vo,	2 00
Hubbard and Kiersted's Water-works Management and Maintenance.....	8vo,	4 00
* Lyndon's Development and Electrical Distribution of Water Power.....	8vo,	3 00
Mason's Water-supply. (Considered Principally from a Sanitary Stand- point.).....	8vo,	4 00
Merriman's Treatise on Hydraulics.....	8vo,	5 00
* Molitor's Hydraulics of Rivers, Weirs and Sluices.....	8vo,	2 00
Morrison and Brodie's High Masonry Dam Design. (In Press.).....		
* Richards's Laboratory Notes on Industrial Water Analysis.....	8vo,	50
Schuyler's Reservoirs for Irrigation, Water-power, and Domestic Water- supply. Second Edition, Revised and Enlarged.....	Large 8vo,	6 00
* Thomas and Watt's Improvement of Rivers.....	4to,	6 00
Turneure and Russell's Public Water-supplies.....	8vo,	5 00
Wegmann's Design and Construction of Dams. 5th Ed., enlarged.....	4to,	6 00
Water-Supply of the City of New York from 1658 to 1895.....	4to,	10 00
Whipple's Value of Pure Water.....	Large 12mo,	1 00
Williams and Hazen's Hydraulic Tables.....	8vo,	1 50
Wilson's Irrigation Engineering.....	8vo,	4 00
Wood's Turbines.....	8vo,	2 50

MATERIALS OF ENGINEERING.

Baker's Roads and Pavements.....	8vo,	5 00
Treatise on Masonry Construction.....	8vo,	5 00
Black's United States Public Works.....	Oblong 4to,	5 00
Blanchard's Bituminous Roads. (In Press.).....		
Bleining's Manufacture of Hydraulic Cement. (In Preparation.).....		
* Bovey's Strength of Materials and Theory of Structures.....	8vo,	7 50
Burr's Elasticity and Resistance of the Materials of Engineering.....	8vo,	7 50
Byrne's Highway Construction.....	8vo,	5 00
Inspection of the Materials and Workmanship Employed in Construction.....	16mo,	3 00
Church's Mechanics of Engineering.....	8vo,	6 00
Du Bois's Mechanics of Engineering.....		
Vol. I. Kinematics, Statics, Kinetics.....	Small 4to,	7 50
Vol. II. The Stresses in Framed Structures, Strength of Materials and Theory of Flexures.....	Small 4to,	10 00
* Eckel's Cements, Limes, and Plasters.....	8vo,	6 00
Stone and Clay Products used in Engineering. (In Preparation.).....		
Fowler's Ordinary Foundations.....	8vo,	3 50
* Greene's Structural Mechanics.....	8vo,	2 50
* Holley's Lead and Zinc Pigments.....	Large 12mo,	3 00
Holley and Ladd's Analysis of Mixed Paints, Color Pigments and Varnishes.....	Large 12mo,	2 50
* Hubbard's Dust Preventives and Road Binders.....	8vo,	3 00
Johnson's (C. M.) Rapid Methods for the Chemical Analysis of Special Steels, Steel-making Alloys and Graphite.....	Large 12mo,	3 00
Johnson's (J. B.) Materials of Construction.....	Large 8vo,	6 00
Keep's Cast Iron.....	8vo,	2 50
Lanza's Applied Mechanics.....	8vo,	7 50
Lowe's Paints for Steel Structures.....	12mo,	1 00

Maire's Modern Pigments and their Vehicles.....	12mo.	\$2 00
Maurer's Technical Mechanics.....	8vo,	4 00
Merrill's Stones for Building and Decoration.....	8vo,	5 00
Merriman's Mechanics of Materials.....	8vo,	5 00
* Strength of Materials.....	12mo,	1 00
Metcalf's Steel. A Manual for Steel-users.....	12mo,	2 00
Morrison's Highway Engineering.....	8vo,	2 50
Patton's Practical Treatise on Foundations.....	8vo,	5 00
Rice's Concrete Block Manufacture.....	8vo,	2 00
Richardson's Modern Asphalt Pavement.....	8vo,	3 00
Richey's Building Foreman's Pocket Book and Ready Reference.....	16mo, mor.	5 00
* Cement Workers' and Plasterers' Edition (Building Mechanics' Ready Reference Series).....	16mo, mor.	1 50
Handbook for Superintendents of Construction.....	16mo, mor.	4 00
* Stone and Brick Masons' Edition (Building Mechanics' Ready Reference Series).....	16mo, mor.	1 50
* Rice's Clays: Their Occurrence, Properties, and Uses.....	8vo,	5 00
* Ries and Leighton's History of the Clay-working Industry of the United States.....	8vo,	2 50
Sabin's Industrial and Artistic Technology of Paint and Varnish.....	8vo,	3 00
* Smith's Strength of Material.....	12mo	1 25
Snow's Principal Species of Wood.....	8vo,	3 50
Spalding's Hydraulic Cement.....	12mo,	2 00
Text-book on Roads and Pavements.....	12mo,	2 00
Taylor and Thompson's Treatise on Concrete, Plain and Reinforced.....	8vo,	5 00
Thurston's Materials of Engineering. In Three Parts.....	8vo,	8 00
Part I. Non-metallic Materials of Engineering and Metallurgy.....	8vo,	2 00
Part II. Iron and Steel.....	8vo,	3 50
Part III. A Treatise on Brasses, Bronzes, and Other Alloys and their Constituents.....	8vo,	2 50
Tillson's Street Pavements and Paving Materials.....	8vo,	4 00
* Trautwine's Concrete, Plain and Reinforced.....	16mo,	2 00
Turneure and Maurer's Principles of Reinforced Concrete Construction. Second Edition, Revised and Enlarged.....	8vo,	3 50
Waterbury's Cement Laboratory Manual.....	12mo,	1 00
Wood's (De V.) Treatise on the Resistance of Materials, and an Appendix on the Preservation of Timber.....	8vo,	2 00
Wood's (M. P.) Rustless Coatings: Corrosion and Electrolysis of Iron and Steel.....	8vo,	4 00

RAILWAY ENGINEERING.

Andrews's Handbook for Street Railway Engineers.....	3×5 inches, mor.	1 25
Berg's Buildings and Structures of American Railroads.....	4to,	5 00
Brooks's Handbook of Street Railroad Location.....	16mc, mor.	1 50
Butts's Civil Engineer's Field-book.....	16mc, mor.	2 50
Crandall's Railway and Other Earthwork Tables.....	8vo,	1 50
Transition Curve.....	16mo, mor.	1 50
* Crockett's Methods for Earthwork Computations.....	8vo,	1 50
Dredge's History of the Pennsylvania Railroad. (1879).....	Paper,	5 00
Fisher's Table of Cubic Yards.....	Cardboard,	25
Godwin's Railroad Engineers' Field-book and Explorers' Guide.....	16mo, mor.	2 50
Hudson's Tables for Calculating the Cubic Contents of Excavations and Embankments.....	8vo,	1 00
Ives and Hilt's Problems in Surveying, Railroad Surveying and Geodesy.....	16mo, mor.	1 50
Molitor and Beard's Manual for Resident Engineers.....	16mo,	1 00
Nagle's Field Manual for Railroad Engineers.....	16mo, mor.	3 00
* Orrock's Railroad Structures and Estimates.....	8vo,	3 00
Philbrick's Field Manual for Engineers.....	16mo, mor.	3 00
Raymond's Railroad Engineering. 3 volumes.		
Vol. I. Railroad Field Geometry. (In Preparation.)		
Vol. II. Elements of Railroad Engineering.....	8vo,	3 50
Vol. III. Railroad Engineer's Field Book. (In Preparation.)		

Roberts' Track Formulæ and Tables. (In Press.)		
Searles's Field Engineering.....	16mo, mor.	\$3 00
Railroad Spiral.....	16mo, mor.	1 50
Taylor's Prismoidal Formulæ and Earthwork.....	8vo,	1 50
* Trautwine's Field Practice of Laying Out Circular Curves for Railroads.	12mo, mor.	2 50
* Method of Calculating the Cubic Contents of Excavations and Embankments by the Aid of Diagrams.....	8vo,	2 00
Webb's Economics of Railroad Construction.....	Large 12mo,	2 50
Railroad Construction.....	16mo, mor.	5 00
Wellington's Economic Theory of the Location of Railways.....	Large 12mo,	5 00
Wilson's Elements of Railroad-Track and Construction.....	12mo,	2 00

DRAWING.

Barr's Kinematics of Machinery.....	8vo,	2 50
* Bartlett's Mechanical Drawing.....	8vo,	3 00
* " " " Abridged Ed.....	8vo,	1 50
Coolidge's Manual of Drawing.....	8vo, paper,	1 00
Coolidge and Freeman's Elements of General Drafting for Mechanical Engineers.....	Oblong 4to,	2 50
Durley's Kinematics of Machines.....	8vo,	4 00
Emch's Introduction to Projective Geometry and its Application.....	8vo,	2 50
French and Ives' Stereotomy.....	8vo,	2 50
Hill's Text-book on Shades and Shadows, and Perspective.....	8vo,	2 00
Jamison's Advanced Mechanical Drawing.....	8vo,	2 00
Elements of Mechanical Drawing.....	8vo,	2 50
Jones's Machine Design:		
Part I. Kinematics of Machinery.....	8vo,	1 50
Part II. Form, Strength, and Proportions of Parts.....	8vo,	3 00
* Kimball and Barr's Machine Design.....	8vo,	3 00
MacCord's Elements of Descriptive Geometry.....	8vo,	3 00
Kinematics; or, Practical Mechanism.....	8vo,	5 00
Mechanical Drawing.....	4to,	4 00
Velocity Diagrams.....	8vo,	1 50
McLeod's Descriptive Geometry.....	Large 12mo,	1 50
* Mahan's Descriptive Geometry and Stone-cutting.....	8vo,	1 50
Industrial Drawing. (Thompson.).....	8vo,	3 50
Moyer's Descriptive Geometry.....	8vo,	2 00
Reed's Topographical Drawing and Sketching.....	4to,	5 00
Reid's Course in Mechanical Drawing.....	8vo,	2 00
Text-book of Mechanical Drawing and Elementary Machine Design.....	8vo,	3 00
Robinson's Principles of Mechanism.....	8vo,	3 00
Schwamb and Merrill's Elements of Mechanism.....	8vo,	3 00
Smith (A. W.) and Marx's Machine Design.....	8vo,	3 00
Smith's (R. S.) Manual of Topographical Drawing. (McMillan.).....	8vo,	2 50
* Titsworth's Elements of Mechanical Drawing.....	Oblong 8vo,	1 25
Warren's Drafting Instruments and Operations.....	12mo,	1 25
Elements of Descriptive Geometry, Shades, and Perspective.....	8vo,	3 50
Elements of Machine Construction and Drawing.....	8vo,	7 50
Elements of Plane and Solid Free-hand Geometrical Drawing....	12mo,	1 00
General Problems of Shades and Shadows.....	8vo,	3 00
Manual of Elementary Problems in the Linear Perspective of Forms and Shadow.....	12mo,	1 00
Manual of Elementary Projection Drawing.....	12mo,	1 50
Plane Problems in Elementary Geometry.....	12mo,	1 25
Weisbach's Kinematics and Power of Transmission. (Hermann and Klein.).....	8vo,	5 00
Wilson's (H. M.) Topographic Surveying.....	8vo,	3 50
* Wilson's (V. T.) Descriptive Geometry.....	8vo,	1 50
Free-hand Lettering.....	8vo,	1 00
Free-hand Perspective.....	8vo,	2 50
Woolf's Elementary Course in Descriptive Geometry.....	Large 8vo,	3 00

ELECTRICITY AND PHYSICS.

* Abegg's Theory of Electrolytic Dissociation. (von Ende.).....	12mo,	\$1 25
Andrews's Hand-book for Street Railway Engineering.....	3×5 inches, mor.	1 25
Anthony and Brackett's Text-book of Physics. (Magie.)....	Large 12mo,	3 00
Anthony and Ball's Lecture-notes on the Theory of Electrical Measurements.....	12mo,	1 00
Benjamin's History of Electricity.....	8vo,	3 00
Voltaic Cell.....	8vo,	3 00
Betts's Lead Refining and Electrolysis.....	8vo,	4 00
Classen's Quantitative Chemical Analysis by Electrolysis. (Boltwood.)	8vo,	3 00
* Collins's Manual of Wireless Telegraphy and Telephony.....	12mo,	1 50
Crehore and Squier's Polarizing Photo-chronograph.....	8vo,	3 00
* Danneel's Electrochemistry. (Merriam.).....	12mo,	1 25
Dawson's "Engineering" and Electric Traction Pocket-book....	16mo, mor.	5 00
Dolezalek's Theory of the Lead Accumulator (Storage Battery). (von Ende.)	12mo,	2 50
Duhem's Thermodynamics and Chemistry. (Burgess.).....	8vo,	4 00
Flather's Dynamometers, and the Measurement of Power.....	12mo,	3 00
* Getman's Introduction to Physical Science.....	12mo,	1 50
Gilbert's De Magnete. (Mottelay).....	8vo,	2 50
* Hanchett's Alternating Currents.....	12mo,	1 00
Hering's Ready Reference Tables (Conversion Factors).....	16mo, mor.	2 50
* Hobart and Ellis's High-speed Dynamo Electric Machinery.....	8vo,	6 00
Holman's Precision of Measurements.....	8vo,	2 00
Telescopic Mirror-scale Method, Adjustments, and Tests....	Large 8vo,	75
* Karapetoff's Experimental Electrical Engineering.....	8vo,	6 00
Kinzbrunner's Testing of Continuous-current Machines.....	8vo,	2 00
Landauer's Spectrum Analysis. (Tingle.).....	8vo,	3 00
Le Chatelier's High-temperature Measurements. (Boudouard—Burgess.)	12mo,	3 00
Löb's Electrochemistry of Organic Compounds. (Lorenz.).....	8vo,	3 00
* Lyndon's Development and Electrical Distribution of Water Power.	8vo,	3 00
* Lyons's Treatise on Electromagnetic Phenomena. Vols. I. and II.	8vo, each,	6 00
* Michie's Elements of Wave Motion Relating to Sound and Light....	8vo,	4 00
Morgan's Outline of the Theory of Solution and its Results.....	12mo,	1 00
* Physical Chemistry for Electrical Engineers.....	12mo,	1 50
* Norris's Introduction to the Study of Electrical Engineering.....	8vo,	2 50
Norris and Dennison's Course of Problems on the Electrical Characteristics of Circuits and Machines. (In Press.)		
* Parshall and Hobart's Electric Machine Design.....	4to, half mor.	12 50
Reagan's Locomotives: Simple, Compound, and Electric. New Edition.		
	Large 12mo,	3 50
* Rosenberg's Electrical Engineering. (Haldane Gee—Kinzbrunner.)	8vo,	2 00
Ryan, Norris, and Hoxie's Electrical Machinery. Vol. I.....	8vo,	2 50
Schapper's Laboratory Guide for Students in Physical Chemistry....	12mo,	1 00
* Tillman's Elementary Lessons in Heat.....	8vo,	1 50
Tory and Pitcher's Manual of Laboratory Physics.....	Large 12mo,	2 00
Ulke's Modern Electrolytic Copper Refining.....	8vo,	3 00

LAW.

* Brennan's Hand-book of Useful Legal Information for Business Men.	16mo, mor.	5 00
* Davis's Elements of Law.....	8vo,	2 50
* Treatise on the Military Law of United States.....	8vo,	7 00
* Dudley's Military Law and the Procedure of Courts-martial..	Large 12mo,	2 50
Manual for Courts-martial.....	16mo, mor.	1 50
Wait's Engineering and Architectural Jurisprudence.....	8vo,	6 00
	Sheep,	6 50
Law of Contracts.....	8vo,	3 00
Law of Operations Preliminary to Construction in Engineering and Architecture.....	8vo,	5 00
	Sheep,	5 50

MATHEMATICS.

Baker's Elliptic Functions.....	8vo,	\$1 50
Briggs's Elements of Plane Analytic Geometry. (Böcher.).....	12mo,	1 00
* Buchanan's Plane and Spherical Trigonometry.....	8vo,	1 00
Byerley's Harmonic Functions.....	8vo,	1 00
Chandler's Elements of the Infinitesimal Calculus.....	12mo,	2 00
* Coffin's Vector Analysis.....	12mo,	2 50
Compton's Manual of Logarithmic Computations.....	12mo,	1 50
* Dickson's College Algebra.....	Large 12mo,	1 50
* Introduction to the Theory of Algebraic Equations.....	Large 12mo,	1 25
Emch's Introduction to Projective Geometry and its Application.....	8vo,	2 50
Fiske's Functions of a Complex Variable.....	8vo,	1 00
Halsted's Elementary Synthetic Geometry.....	8vo,	1 50
Elements of Geometry.....	8vo,	1 75
* Rational Geometry.....	12mo,	1 50
Synthetic Projective Geometry.....	8vo,	1 00
Hancock's Lectures on the Theory of Elliptic Functions. (In Press.).....	8vo,	1 00
Hyde's Grassmann's Space Analysis.....	8vo,	1 00
* Johnson's (J. B.) Three-place Logarithmic Tables: Vest-pocket size, paper,		15
* 100 copies,		5 00
* Mounted on heavy cardboard, 8×10 inches,		25
* 10 copies,		2 00
Johnson's (W. W.) Abridged Editions of Differential and Integral Calculus.		
Large 12mo, 1 vol.		2 50
Curve Tracing in Cartesian Co-ordinates.....	12mo,	1 00
Differential Equations.....	8vo,	1 00
Elementary Treatise on Differential Calculus.....	Large 12mo,	1 50
Elementary Treatise on the Integral Calculus.....	Large 12mo,	1 50
* Theoretical Mechanics.....	12mo,	3 00
Theory of Errors and the Method of Least Squares.....	12mo,	1 50
Treatise on Differential Calculus.....	Large 12mo,	3 00
Treatise on the Integral Calculus.....	Large 12mo,	3 00
Treatise on Ordinary and Partial Differential Equations.....	Large 12mo,	3 50
Karapetoff's Engineering Applications of Higher Mathematics. (In Preparation.)		
Laplace's Philosophical Essay on Probabilities. (Truscott and Emory.)	12mo,	2 00
* Ludlow and Bass's Elements of Trigonometry and Logarithmic and Other		
Tables.....	8vo,	3 00
* Trigonometry and Tables published separately.....	Each,	2 00
* Ludlow's Logarithmic and Trigonometric Tables.....	8vo,	1 00
Macfarlane's Vector Analysis and Quaternions.....	8vo,	1 00
McMahon's Hyperbolic Functions.....	8vo,	1 00
Manning's Irrational Numbers and their Representation by Sequences and		
Series.....	12mo,	1 25
Mathematical Monographs. Edited by Mansfield Merriman and Robert		
S. Woodward.....	Octavo, each	1 00
No. 1. History of Modern Mathematics, by David Eugene Smith.		
No. 2. Synthetic Projective Geometry, by George Bruce Halsted.		
No. 3. Determinants, by Laenas Gifford Weld. No. 4. Hyper-		
bolic Functions, by James McMahon. No. 5. Harmonic Func-		
tions, by William E. Byerly. No. 6. Grassmann's Space Analysis,		
by Edward W. Hyde. No. 7. Probability and Theory of Errors,		
by Robert S. Woodward. No. 8. Vector Analysis and Quaternions,		
by Alexander Macfarlane. No. 9. Differential Equations, by		
William Woolsey Johnson. No. 10. The Solution of Equations,		
by Mansfield Merriman. No. 11. Functions of a Complex Variable,		
by Thomas S. Fiske.		
Maurer's Technical Mechanics.....	8vo,	4 00
Merriman's Method of Least Squares.....	8vo,	2 00
Solution of Equations.....	8vo,	1 00
Rice and Johnson's Differential and Integral Calculus. 2 vols. in one.		
Large 12mo,		1 50
Elementary Treatise on the Differential Calculus.....	Large 12mo,	3 00
Smith's History of Modern Mathematics.....	8vo,	1 00
* Veblen and Lennes's Introduction to the Real Infinitesimal Analysis of One		
Variable.....	8vo,	2 00

* Waterbury's Vest Pocket Hand-book of Mathematics for Engineers.

	2 $\frac{1}{4}$ × 5 $\frac{3}{8}$ inches, mor.	\$1 00
* Enlarged Edition, Including Tables	mor.	1 50
Weld's Determinants.....	8vo,	1 00
Wood's Elements of Co-ordinate Geometry.....	8vo,	2 00
Woodward's Probability and Theory of Errors.....	8vo,	1 00

MECHANICAL ENGINEERING.

MATERIALS OF ENGINEERING, STEAM-ENGINES AND BOILERS.

Bacon's Forge Practice.....	12mo,	1 50
Baldwin's Steam Heating for Buildings.....	12mo,	2 50
Barr's Kinematics of Machinery.....	8vo,	2 50
* Bartlett's Mechanical Drawing.....	8vo,	3 00
* " " " Abridged Ed.....	8vo,	1 50
* Burr's Ancient and Modern Engineering and the Isthmian Canal.....	8vo,	3 50
Carpenter's Experimental Engineering.....	8vo,	6 00
Heating and Ventilating Buildings.....	8vo,	4 00
* Clerk's The Gas, Petrol and Oil Engine.....	8vo,	4 00
Compton's First Lessons in Metal Working.....	12mo,	1 50
Compton and De Groodt's Speed Lathe.....	12mo,	1 50
Coolidge's Manual of Drawing.....	8vo, paper,	1 00
Coolidge and Freeman's Elements of General Drafting for Mechanical Engineers.....	Oblong 4to,	2 50
Cromwell's Treatise on Belts and Pulleys.....	12mo,	1 50
Treatise on Toothed Gearing.....	12mo,	1 50
Dingey's Machinery Pattern Making.....	12mo,	2 00
Durley's Kinematics of Machines.....	8vo,	4 00
Flanders's Gear-cutting Machinery.....	Large 12mo,	3 00
Flather's Dynamometers and the Measurement of Power.....	12mo,	3 00
Rope Driving.....	12mo,	2 00
Gill's Gas and Fuel Analysis for Engineers.....	12mo,	1 25
Goss's Locomotive Sparks.....	8vo,	2 00
Greene's Pumping Machinery. (In Preparation.).....		
Hering's Ready Reference Tables (Conversion Factors).....	16mo, mor.	2 50
* Hobart and Ellis's High Speed Dynamo Electric Machinery.....	8vo,	6 00
Hutton's Gas Engine.....	8vo,	5 00
Jamison's Advanced Mechanical Drawing.....	8vo,	2 00
Elements of Mechanical Drawing.....	8vo,	2 50
Jones's Gas Engine.....	8vo,	4 00
Machine Design:		
Part I. Kinematics of Machinery.....	8vo,	1 50
Part II. Form, Strength, and Proportions of Parts.....	8vo,	3 00
Kent's Mechanical Engineer's Pocket-Book.....	16mo, mor.	5 00
Kerr's Power and Power Transmission.....	8vo,	2 00
* Kimball and Barr's Machine Design.....	8vo,	3 00
* Levin's Gas Engine.....	8vo,	4 00
Leonard's Machine Shop Tools and Methods.....	8vo,	4 00
* Lorenz's Modern Refrigerating Machinery. (Pope, Haven, and Dean).....	8vo,	4 00
MacCord's Kinematics; or, Practical Mechanism.....	8vo,	5 00
Mechanical Drawing.....	4to,	4 00
Velocity Diagrams.....	8vo,	1 50
MacFarland's Standard Reduction Factors for Gases.....	8vo,	1 50
Mahan's Industrial Drawing. (Thompson.).....	8vo,	3 50
Mehrtens's Gas Engine Theory and Design.....	Large 12mo,	2 50
Oberg's Handbook of Small Tools.....	Large 12mo,	3 00
* Parshall and Hobart's Electric Machine Design. Small 4to, half leather,		12 50
Peele's Compressed Air Plant for Mines.....	8vo,	3 00
Poole's Calorific Power of Fuels.....	8vo,	3 00
* Porter's Engineering Reminiscences, 1855 to 1882.....	8vo,	3 00
Reid's Course in Mechanical Drawing.....	8vo,	2 00
Text-book of Mechanical Drawing and Elementary Machine Design.....	8vo,	3 00

Richards's Compressed Air.	12mo,	\$1 50
Robinson's Principles of Mechanism.	8vo,	3 00
Schwamb and Merrill's Elements of Mechanism.	8vo,	3 00
Smith (A. W.) and Marx's Machine Design.	8vo,	3 00
Smith's (O.) Press-working of Metals.	8vo,	3 00
Sorel's Carbureting and Combustion in Alcohol Engines. (Woodward and Preston.)	Large 12mo,	3 00
Stone's Practical Testing of Gas and Gas Meters.	8vo,	3 50
Thurston's Animal as a Machine and Prime Motor, and the Laws of Energetics.	12mo,	1 00
Treatise on Friction and Lost Work in Machinery and Mill Work.	8vo,	3 00
* Tillson's Complete Automobile Instructor.	16mo,	1 50
* Tittsworth's Elements of Mechanical Drawing.	Oblong 8vo,	1 25
Warren's Elements of Machine Construction and Drawing.	8vo,	7 50
* Waterbury's Vest Pocket Hand-book of Mathematics for Engineers.	2 $\frac{1}{8}$ × 5 $\frac{3}{8}$ inches, mor.	1 00
* Enlarged Edition, Including Tables.	mor.	1 50
Weissbach's Kinematics and the Power of Transmission. (Herrmann—Klein.)	8vo,	5 00
Machinery of Transmission and Governors. (Herrmann—Klein.)	8vo,	5 00
Wood's Turbines.	8vo,	2 50

MATERIALS OF ENGINEERING.

* Bovey's Strength of Materials and Theory of Structures.	8vo,	7 50
Burr's Elasticity and Resistance of the Materials of Engineering.	8vo,	7 50
Church's Mechanics of Engineering.	8vo,	6 00
* Greene's Structural Mechanics.	8vo,	2 50
* Holley's Lead and Zinc Pigments.	Large 12mo	3 00
Holley and Ladd's Analysis of Mixed Paints, Color Pigments, and Varnishes.	Large 12mo,	2 50
Johnson's (C. M.) Rapid Methods for the Chemical Analysis of Special Steels, Steel-Making Alloys and Graphite.	Large 12mo,	3 00
Johnson's (J. B.) Materials of Construction.	8vo,	6 00
Keep's Cast Iron.	8vo,	2 50
Lanza's Applied Mechanics.	8vo,	7 50
Maire's Modern Pigments and their Vehicles.	12mo,	2 00
Maurer's Technical Mechanics.	8vo,	4 00
Merriman's Mechanics of Materials.	8vo,	5 00
* Strength of Materials.	12mo,	1 00
Metcalf's Steel. A Manual for Steel-users.	12mo,	2 00
Sabin's Industrial and Artistic Technology of Paint and Varnish.	8vo,	3 00
Smith's ((A. W.) Materials of Machines.	12mo,	1 00
* Smith's (H. E.) Strength of Material.	12mo,	1 25
Thurston's Materials of Engineering.	3 vols., 8vo,	8 00
Part I. Non-metallic Materials of Engineering.	8vo,	2 00
Part II. Iron and Steel.	8vo,	3 50
Part III. A Treatise on Brasses, Bronzes, and Other Alloys and their Constituents.	8vo,	2 50
Wood's (De V.) Elements of Analytical Mechanics.	8vo,	3 00
Treatise on the Resistance of Materials and an Appendix on the Preservation of Timber.	8vo,	2 00
Wood's (M. P.) Rustless Coatings: Corrosion and Electrolysis of Iron and Steel.	8vo,	4 00

STEAM-ENGINES AND BOILERS.

Berry's Temperature-entropy Diagram.	12mo,	2 00
Carnot's Reflections on the Motive Power of Heat. (Thurston.)	12mo,	1 50
Chase's Art of Pattern Making.	12mo,	2 50

Creighton's Steam-engine and other Heat Motors.	8vo,	\$5 00
Dawson's "Engineering" and Electric Traction Pocket-book. ...	16mo, mor.	5 00
* Gebhardt's Steam Power Plant Engineering.	8vo,	6 00
Goss's Locomotive Performance.	8vo,	5 00
Hemenway's Indicator Practice and Steam-engine Economy.	12mo,	2 00
Hutton's Heat and Heat-engines.	8vo,	5 00
Mechanical Engineering of Power Plants.	8vo,	5 00
Kent's Steam boiler Economy.	8vo,	4 00
Kneass's Practice and Theory of the Injector.	8vo,	1 50
MacCord's Slide-valves.	8vo,	2 00
Meyer's Modern Locomotive Construction.	4to,	10 00
Moyer's Steam Turbine.	8vo,	4 00
Peabody's Manual of the Steam-engine Indicator.	12mo,	1 50
Tables of the Properties of Steam and Other Vapors and Temperature-Entropy Table.	8vo,	1 00
Thermodynamics of the Steam-engine and Other Heat-engines. ...	8vo,	5 00
Valve-gears for Steam-engines.	8vo,	2 50
Peabody and Miller's Steam-boilers.	8vo,	4 00
Pupin's Thermodynamics of Reversible Cycles in Gases and Saturated Vapors. (Osterberg.).	12mo,	1 25
Reagan's Locomotives: Simple, Compound, and Electric. New Edition. Large 12mo,		3 50
Sinclair's Locomotive Engine Running and Management.	12mo,	2 00
Smart's Handbook of Engineering Laboratory Practice.	12mo,	2 50
Snow's Steam-boiler Practice.	8vo,	3 00
Spangler's Notes on Thermodynamics.	12mo,	1 00
Valve-gears.	8vo,	2 50
Spangler, Greene, and Marshall's Elements of Steam-engineering. ...	8vo,	3 00
Thomas's Steam-turbines.	8vo,	4 00
Thurston's Handbook of Engine and Boiler Trials, and the Use of the Indicator and the Prony Brake.	8vo,	5 00
Handy Tables.	8vo,	1 50
Manual of Steam-boilers, their Designs, Construction, and Operation	8vo,	5 00
Manual of the Steam-engine. 2 vols.,	8vo,	10 00
Part I. History, Structure, and Theory	8vo,	6 00
Part II. Design, Construction, and Operation.	8vo,	6 00
Wehrenfennig's Analysis and Softening of Boiler Feed-water. (Patterson.)	8vo,	4 00
Weisbach's Heat, Steam, and Steam-engines. (Du Bois.).	8vo,	5 00
Whitham's Steam-engine Design.	8vo,	5 00
Wood's Thermodynamics, Heat Motors, and Refrigerating Machines. .	8vo,	4 00

MECHANICS PURE AND APPLIED.

Church's Mechanics of Engineering.	8vo,	6 00
Notes and Examples in Mechanics.	8vo,	2 00
Dana's Text-book of Elementary Mechanics for Colleges and Schools .	12mo,	1 50
Du Bois's Elementary Principles of Mechanics:		
Vol. I. Kinematics.	8vo,	3 50
Vol. II. Statics.	8vo,	4 00
Mechanics of Engineering. Vol. I.	Small 4to,	7 50
Vol. II.	Small 4to,	10 00
* Greene's Structural Mechanics.	8vo,	2 50
Hartmann's Elementary Mechanics for Engineering Students. (In Press.)		
James's Kinematics of a Point and the Rational Mechanics of a Particle. Large 12mo,		2 00
* Johnson's (W. W.) Theoretical Mechanics.	12mo,	3 00
Lanza's Applied Mechanics.	8vo,	7 50
* Martin's Text Book on Mechanics, Vol. I, Statics.	12mo,	1 25
* Vol. II, Kinematics and Kinetics.	12mo,	1 50
Maurer's Technical Mechanics.	8vo,	4 00
* Merriman's Elements of Mechanics.	12mo,	1 00
Mechanics of Materials.	8vo,	5 00
* Michie's Elements of Analytical Mechanics.	8vo,	4 00

Robinson's Principles of Mechanism.	8vo,	\$3 00
Sanborn's Mechanics Problems.	Large 12mo,	1 50
Schwamb and Merrill's Elements of Mechanism.	8vo,	3 00
Wood's Elements of Analytical Mechanics.	8vo,	3 00
Principles of Elementary Mechanics.	12mo,	1 25

MEDICAL.

* Abderhalden's Physiological Chemistry in Thirty Lectures. (Hall and Defren.)	8vo,	5 00
von Behring's Suppression of Tuberculosis. (Bolduan.)	12mo,	1 00
Bolduan's Immune Sera.	12mo,	1 50
Bordet's Studies in Immunity. (Gay.)	8vo,	6 00
Chapin's The Sources and Modes of Infection. (In Press.)		
Davenport's Statistical Methods with Special Reference to Biological Variations.	16mo, mor.	1 50
Ehrlich's Collected Studies on Immunity. (Bolduan.)	8vo,	6 00
* Fischer's Physiology of Alimentation.	Large 12mo,	2 00
de Fursac's Manual of Psychiatry. (Rosanoff and Collins.)	Large 12mo,	2 50
Hammarsten's Text-book on Physiological Chemistry. (Mandel.)	8vo,	4 00
Jackson's Directions for Laboratory Work in Physiological Chemistry.	8vo,	1 25
Lassar-Cohn's Practical Urinary Analysis. (Lorenz.)	12mo,	1 00
Mandel's Hand-book for the Bio-Chemical Laboratory.	12mo,	1 50
* Nelson's Analysis of Drugs and Medicines.	12mo,	3 00
* Pauli's Physical Chemistry in the Service of Medicine. (Fischer.)	12mo,	1 25
* Pozzi-Escot's Toxins and Venoms and their Antibodies. (Cohn.)	12mo,	1 00
Rostoski's Serum Diagnosis. (Bolduan.)	12mo,	1 00
Ruddiman's Incompatibilities in Prescriptions.	8vo,	2 00
Whys in Pharmacy.	12mo,	1 00
Salkowski's Physiological and Pathological Chemistry. (Orndorff.)	8vo,	2 50
* Satterlee's Outlines of Human Embryology.	12mo,	1 25
Smith's Lecture Notes on Chemistry for Dental Students.	8vo,	2 50
* Whipple's Typhoid Fever.	Large 12mo,	3 00
* Woodhull's Military Hygiene for Officers of the Line.	Large 12mo,	1 50
* Personal Hygiene.	12mo,	1 00
Worcester and Atkinson's Small Hospitals Establishment and Maintenance, and Suggestions for Hospital Architecture, with Plans for a Small Hospital.	12mo,	1 25

METALLURGY.

Betts's Lead Refining by Electrolysis.	8vo,	4 00
Bolland's Encyclopedia of Founding and Dictionary of Foundry Terms used in the Practice of Moulding.	12mo,	3 00
Iron Founder.	12mo,	2 50
" Supplement.	12mo,	2 50
Douglas's Untechnical Addresses on Technical Subjects.	12mo,	1 00
Goesel's Minerals and Metals: A Reference Book.	16mo, mor.	3 00
* Iles's Lead-smelting.	12mo,	2 50
Johnson's Rapid Methods for the Chemical Analysis of Special Steels, Steel-making Alloys and Graphite.	Large 12mo,	3 00
Keep's Cast Iron.	8vo,	2 50
Le Chatelier's High-temperature Measurements. (Boudouard—Burgess.)	12mo,	3 00
Metcalf's Steel. A Manual for Steel-users.	12mo,	2 00
Minet's Production of Aluminum and its Industrial Use. (Waldo.)	12mo,	2 50
* Ruer's Elements of Metallography. (Mathewson.)	8vo,	3 00
Smith's Materials of Machines.	12mo,	1 00
Tate and Stone's Foundry Practice.	12mo,	2 00
Thurston's Materials of Engineering. In Three Parts.	8vo,	8 00
Part I. Non-metallic Materials of Engineering, see Civil Engineering, page 9.		
Part II. Iron and Steel.	8vo,	3 50
Part III. A Treatise on Brasses, Bronzes, and Other Alloys and their Constituents.	8vo,	2 50

Ulke's Modern Electrolytic Copper Refining.....	8vo,	\$3 00
West's American Foundry Practice.....	12mo,	£ 50
Moulders' Text Book.....	12mo,	2 50

MINERALOGY.

Baskerville's Chemical Elements. (In Preparation.)		
* Browning's Introduction to the Rarer Elements.....	8vo,	1 50
Brush's Manual of Determinative Mineralogy. (Penfield.).....	8vo,	4 00
Butler's Pocket Hand-book of Minerals.....	16mo, mor,	3 00
Chester's Catalogue of Minerals.....	8vo, paper,	1 00
	Cloth,	1 25
* Crane's Gold and Silver.....	8vo,	5 00
Dana's First Appendix to Dana's New "System of Mineralogy".....	Large 8vo,	1 00
Dana's Second Appendix to Dana's New "System of Mineralogy."		
	Large 8vo,	1 50
Manual of Mineralogy and Petrography.....	12mo,	2 00
Minerals and How to Study Them.....	12mo,	1 50
System of Mineralogy.....	Large 8vo, half leather,	12 50
Text-book of Mineralogy.....	8vo,	4 00
Douglas's Untechnical Addresses on Technical Subjects.....	12mo,	1 00
Eakle's Mineral Tables.....	8vo,	1 25
Eckel's Stone and Clay Products Used in Engineering. (In Preparation.)		
Goesel's Minerals and Metals: A Reference Book.....	16mo, mor.	3 00
Groth's The Optical Properties of Crystals. (Jackson.) (In Press.)		
Groth's Introduction to Chemical Crystallography (Marshall).....	12mo,	1 25
* Hayes's Handbook for Field Geologists.....	16mo, mor.	1 50
Iddings's Igneous Rocks.....	8vo,	5 00
Rock Minerals.....	8vo,	5 00
Johannsen's Determination of Rock-forming Minerals in Thin Sections.....	8vo,	
	With Thumb Index	5 00
* Martin's Laboratory Guide to Qualitative Analysis with the Blow-pipe.....	12mo,	60
Merrill's Non-metallic Minerals: Their Occurrence and Uses.....	8vo,	4 00
Stones for Building and Decoration.....	8vo,	5 00
* Penfield's Notes on Determinative Mineralogy and Record of Mineral Tests.....	8vo, paper,	50
Tables of Minerals, Including the Use of Minerals and Statistics of Domestic Production.....	8vo,	1 00
* Pirsson's Rocks and Rock Minerals.....	12mo,	2 50
* Richards's Synopsis of Mineral Characters.....	12mo, mor.	1 25
* Ries's Clays: Their Occurrence, Properties and Uses.....	8vo,	5 00
* Ries and Leighton's History of the Clay-working Industry of the United States.....	8vo,	2 50
* Tillman's Text-book of Important Minerals and Rocks.....	8vo,	2 00
Washington's Manual of the Chemical Analysis of Rocks.....	8vo,	2 00

MINING.

* Beard's Mine Gases and Explosions.....	Large 12mo,	3 00
* Crane's Gold and Silver.....	8vo,	5 00
* Index of Mining Engineering Literature.....	8vo,	4 00
	* 8vo, mor.	5 00
Ore Mining Methods. (In Press.)		
Douglas's Untechnical Addresses on Technical Subjects.....	12mo,	1 00
Eissler's Modern High Explosives.....	8vo,	4 00
Goesel's Minerals and Metals: A Reference Book.....	16mo, mor.	3 00
Ihlseng's Manual of Mining.....	8vo,	5 00
* Iles's Lead Smelting.....	12mo,	2 50
Peele's Compressed Air Plant for Mines.....	8vo,	3 00
Riemer's Shaft Sinking Under Difficult Conditions. (Corning and Peele.)	8vo,	3 00
* Weaver's Military Explosives.....	8vo,	3 00
Wilson's Hydraulic and Placer Mining. 2d edition, rewritten.....	12mo,	2 50
Treatise on Practical and Theoretical Mine Ventilation.....	12mo,	1 25

SANITARY SCIENCE.

Association of State and National Food and Dairy Departments, Hartford Meeting, 1906.	8vo,	\$3 00
Jamestown Meeting, 1907.	8vo,	3 00
* Bashore's Outlines of Practical Sanitation.	12mo,	1 25
Sanitation of a Country House.	12mo,	1 00
Sanitation of Recreation Camps and Parks.	12mo,	1 00
Chapin's The Sources and Modes of Infection. (In Press.)		
Folwell's Sewerage. (Designing, Construction, and Maintenance.)....	8vo,	3 00
Water-supply Engineering.	8vo,	4 00
Fowler's Sewage Works Analyses.	12mo,	2 00
Fuertes's Water-filtration Works.	12mo,	2 50
Water and Public Health.	12mo,	1 50
Gerhard's Guide to Sanitary Inspections.	12mo,	1 50
* Modern Baths and Bath Houses.	8vo,	3 00
Sanitation of Public Buildings.	12mo,	1 50
* The Water Supply, Sewerage, and Plumbing of Modern City Buildings.	8vo,	4 00
Hazen's Clean Water and How to Get It.	Large 12mo,	1 50
Filtration of Public Water-supplies.	8vo,	3 00
Kinnicut, Winslow and Pratt's Purification of Sewage. (In Preparation.)		
Leach's Inspection and Analysis of Food with Special Reference to State Control.	8vo,	7 50
Mason's Examination of Water. (Chemical and Bacteriological).	12mo,	1 25
Water-supply. (Considered principally from a Sanitary Standpoint).	8vo,	4 00
* Merriman's Elements of Sanitary Engineering.	8vo,	2 00
Ogden's Sewer Construction.	8vo,	3 00
Sewer Design.	12mo,	2 00
Parsons's Disposal of Municipal Refuse.	8vo,	2 00
Prescott and Winslow's Elements of Water Bacteriology, with Special Reference to Sanitary Water Analysis.	12mo,	1 50
* Price's Handbook on Sanitation.	12mo,	1 50
Richards's Cost of Cleanness.	12mo,	1 00
Cost of Food. A Study in Diaries.	12mo,	1 00
Cost of Living as Modified by Sanitary Science.	12mo,	1 00
Cost of Shelter.	12mo,	1 00
* Richards and Williams's Dietary Computer.	8vo,	1 50
Richards and Woodman's Air, Water, and Food from a Sanitary Standpoint.	8vo,	2 00
* Richey's Plumbers', Steam-fitters', and Tinnners' Edition (Building Mechanics' Ready Reference Series).	16mo, mor,	1 50
Rideal's Disinfection and the Preservation of Food.	8vo,	4 00
Sewage and Bacterial Purification of Sewage.	8vo,	4 00
Soper's Air and Ventilation of Subways.	12mo,	2 50
Turneaure and Russell's Public Water-supplies.	8vo,	5 00
Venable's Garbage Crematories in America.	8vo,	2 00
Method and Devices for Bacterial Treatment of Sewage.	8vo,	3 00
Ward and Whipple's Freshwater Biology. (In Press.)		
Whipple's Microscopy of Drinking-water.	8vo,	3 50
* Typhoid Fever.	Large 12mo,	3 00
Value of Pure Water.	Large 12mo,	1 00
Winslow's Systematic Relationship of the Coccaceæ.	Large 12mo,	2 50

MISCELLANEOUS.

Emmons's Geological Guide-book of the Rocky Mountain Excursion of the International Congress of Geologists.	Large 8vo,	1 50
Ferrel's Popular Treatise on the Winds.	8vo,	4 00
Fitzgerald's Boston Machinist.	18mo,	1 00
Gannett's Statistical Abstract of the World.	24mo,	75
Haines's American Railway Management.	12mo,	2 50
Hanausek's The Microscopy of Technical Products. (Winton)	8vo,	5 00

Jacobs's Betterment Briefs. A Collection of Published Papers on Organized Industrial Efficiency.	8vo,	\$3 50
Metcalfe's Cost of Manufactures, and the Administration of Workshops.	8vo,	5 00
Putnam's Nautical Charts.	8vo,	2 00
Ricketts's History of Rensselaer Polytechnic Institute 1824-1894.		
	Large 12mo,	3 00
Rotherham's Emphasised New Testament.	Large 8vo,	2 00
Rust's Ex-Meridian Altitude, Azimuth and Star-finding Tables.	8vo,	5 00
Standage's Decoration of Wood, Glass, Metal, etc.	12mo,	2 00
Thome's Structural and Physiological Botany. (Bennett).....	16mo,	2 25
Westermaier's Compendium of General Botany. (Schneider).....	8vo,	2 00
Winslow's Elements of Applied Microscopy.	12mo,	1 50

HEBREW AND CHALDEE TEXT-BOOKS.

Gesenius's Hebrew and Chaldee Lexicon to the Old Testament Scriptures. (Tregelles.)	Small 4to, half mor,	5 00
Green's Elementary Hebrew Grammar.	12mo,	1 25

UNIVERSITY OF ILLINOIS-URBANA

512.94M55S1906

C002

THE SOLUTION OF EQUATIONS 4TH ED., ENL.



3 0112 017082881